

## Accepted Manuscript

Title: Integrating psychological and neurobiological considerations regarding the development and maintenance of specific Internet-use disorders: An Interaction of Person-Affect-Cognition-Execution (I-PACE) model

Author: Matthias Brand Kimberly Young Christian Laier  
Klaus Wölfling Marc N. Potenza



PII: S0149-7634(16)30262-7  
DOI: <http://dx.doi.org/doi:10.1016/j.neubiorev.2016.08.033>  
Reference: NBR 2579

To appear in:

Received date: 29-4-2016  
Revised date: 11-8-2016  
Accepted date: 27-8-2016

Please cite this article as: Brand, Matthias, Young, Kimberly, Laier, Christian, Wölfling, Klaus, Potenza, Marc N., Integrating psychological and neurobiological considerations regarding the development and maintenance of specific Internet-use disorders: An Interaction of Person-Affect-Cognition-Execution (I-PACE) model. *Neuroscience and Biobehavioral Reviews* <http://dx.doi.org/10.1016/j.neubiorev.2016.08.033>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Integrating psychological and neurobiological considerations regarding the development and maintenance of specific Internet-use disorders: An Interaction of Person-Affect-Cognition-Execution (I-PACE) model**

*Matthias Brand<sup>1,2</sup>, Kimberly Young<sup>3</sup>, Christian Laier<sup>1</sup>, Klaus Wölfling<sup>4</sup>, & Marc N. Potenza<sup>5,6,7</sup>*

<sup>1</sup> General Psychology: Cognition and Center for Behavioral Addiction Research (CeBAR), University of Duisburg-Essen, Germany

<sup>2</sup> Erwin L. Hahn Institute for Magnetic Resonance Imaging, Essen, Germany

<sup>3</sup> Center for Internet Addiction, Russell J. Jandoli School of Journalism and Mass Communication, St. Bonaventure University, Olean, USA

<sup>4</sup> Outpatient Clinic for Behavioral Addiction, Department of Psychosomatic Medicine and Psychotherapy, University Medical Center, Johannes Gutenberg University Mainz, Germany

<sup>5</sup> Departments of Psychiatry, Neuroscience and Child Study, Yale University School of Medicine, New Haven, USA

<sup>6</sup> National Center on Addiction and Substance Abuse, Yale University School of Medicine, New Haven, USA

<sup>7</sup> Connecticut Mental Health Center, New Haven, USA

**Corresponding author:**

***Prof. Dr. Matthias Brand***

General Psychology: Cognition  
and Center for Behavioral Addiction Research  
and Erwin L. Hahn Institute for Magnetic Resonance Imaging  
University of Duisburg-Essen  
Forsthausweg 2, 47057 Duisburg  
Tel: ++49-203-3792541 Fax: ++49-203-3791846  
E-mail: matthias.brand@uni-due.de

## Highlights

- Internet-use disorders involve interactions of Person-Affect-Cognition-Execution
- Affect and cognition mediate Internet-use-disorder processes
- Executive and affective features interact in Internet-use-disorder processes
- Conditioning processes impact interactions of Person-Affect-Cognition-Execution
- Ventral striatal and prefrontal function is linked to Internet-use disorders

## Abstract

Within the last two decades, many studies have addressed the clinical phenomenon of Internet-use disorders, with a particular focus on Internet-gaming disorder. Based on previous theoretical considerations and empirical findings, we suggest an Interaction of Person-Affect-Cognition-Execution (I-PACE) model of specific Internet-use disorders. The I-PACE model is a theoretical framework for the processes underlying the development and maintenance of an addictive use of certain Internet applications or sites promoting gaming, gambling, pornography viewing, shopping, or communication. The model is composed as a process model. Specific Internet-use disorders are considered to be the consequence of interactions between predisposing factors, such as neurobiological and psychological constitutions, moderators, such as coping styles and Internet-related cognitive biases, and mediators, such as affective and cognitive responses to situational triggers in combination with reduced executive functioning. Conditioning processes may strengthen these associations within an addiction process. Although the hypotheses regarding the mechanisms underlying the development and maintenance of specific Internet-use disorders, summarized in the I-PACE model, must be further tested empirically, implications for treatment interventions are suggested.

**Keywords:** Internet addiction, Internet-gaming disorder, Internet-use disorders, Cue-reactivity, Inhibitory control, Executive functions

## 1 Introduction

The first description of a patient with symptoms of Internet addiction was published by Kimberly Young (1996). Although the extent to which the Internet may be considered as a focus of an addiction or facilitates addictive behaviors (or whether excessive or problematic engagement in Internet-related behaviors should be considered within an addiction framework at all) remains debated (Petry & O'Brian, 2013), there have been significant changes in the availability and use of the Internet since 1996. Within the last two decades, research on Internet addiction has grown significantly. Many studies have assessed epidemiological factors of an addictive use of the Internet, its prevalence in different countries as well as its comorbidities and personality correlates (see recent reviews by Cash, Rae, Steel, & Winkler, 2012; Kuss & Lopez-Fernandez, 2016; Pezoa-Jares, Espinoza-Luna, & Vasquez-Medina, 2012; Pontes, Kuss, & Griffiths, 2015; Spada, 2014; Suissa, 2015). In the fifth edition of the Diagnostic and Statistical Manual (DSM-5) (APA, 2013), Internet-gaming disorder - as one specific type of Internet addiction - has recently been included in section III emphasizing that it is most likely that this condition has clinical significance, but that more research is needed to ensure its clinical relevance and the exact phenomenology. Although the term Internet addiction is discussed controversially (Starcevic, 2013), it is arguably the most frequently used term in international publications (e.g., Brand, Laier, & Young, 2014; Brand, Young, & Laier, 2014; Chou, Condron, & Belland, 2005; Dong, Shen, Huang, & Du, 2013; Douglas et al., 2008; Griffiths, 1999; Hansen, 2002; Kuss & Griffiths, 2011a; Kuss, Griffiths, Karila, & Billieux, 2014; Weinstein & Lejoyeux, 2010; Widyanto & Griffiths, 2006; Young, 1998, 2004; Young, Yue, & Ying, 2011). However, given controversies regarding the usage of the term addiction and to be consistent with existing nomenclature in DSM-5 and proposed nomenclature in ICD-11, we will more

frequently use the term Internet-use disorder(s) except when it is more precise to use the term Internet addiction (e.g., when referring to prior literature).

Although the DSM-5 focuses on Internet gaming, a meaningful number of authors indicate that treatment-seeking individuals may also use other Internet applications or sites addictively. Prominent examples include gambling, pornography, social-networking, and shopping sites (Brand, Young, et al., 2014; Griffiths, 2012; Kuss & Griffiths, 2011b; A. Müller, Brand, Mitchell, & de Zwaan, in press; K. W. Müller et al., 2016; Young, Pistner, O'Mara, & Buchanan, 1999). The Internet activity of individuals reporting features of addictive usage should therefore be specified, because individuals are not addicted to the medium itself *per se*, but to the content they are using (see comprehensive discussion in Starcevic, 2013). Empirical evidence also suggests differentiating between a more generalized Internet addiction and specific types of addictive Internet usage (e.g., Montag et al., 2015; Pawlikowski et al., 2014). Consistent with this notion, we argue to use the term specific Internet-use disorders, which implies that the content used should be specified, for example Internet-gaming disorder, Internet-gambling disorder, Internet-pornography-viewing disorder, etc. (Brand, Young, et al., 2014). An awareness of the common and distinct processes behind these phenomena may have a significant impact on policies, prevention efforts and clinical treatments.

For both research and clinical practice, theoretical models of the mechanisms underlying the development and maintenance of addictive behaviors are very important. For Internet addiction, two theoretical models have been published in 2014, one by Brand, Young, et al. (2014), and another, which focuses on Internet gaming, by Dong and Potenza (2014). Since the publication of these two models, new research findings exist, which partly confirm certain theoretical assumptions of the models, but which also give rise to new ideas on the mechanisms involved in the

addiction process. Consequently, we think that revising our model on specific Internet-use disorders (Brand, Young, et al., 2014) is timely, given that theoretical models and frameworks should be modified based on new data emerging from research.

The aim of the current article is to suggest a revised version of our model on specific Internet-use disorders. The specific goals are as follows. First, we integrate current research on Internet-use disorders into the theoretical model. We also integrate findings and theoretical assumptions from other research areas, e.g., by referring to concepts known from substance-dependence research. This is consistent with the idea to classify Internet-use disorders and other behavioral addictions together with substance-use disorders as addictive behaviors (cf. Chamberlain et al., 2015; Derbyshire & Grant, 2015; Fauth-Bühler & Mann, 2015; Fauth-Bühler, Mann, & Potenza, 2016; Grant, Brewer, & Potenza, 2006; Grant & Chamberlain, 2015; Grant, Potenza, Weinstein, & Gorelick, 2010; Kraus, Voon, & Potenza, 2016; Potenza, 2006; Robbins & Clark, 2015). Second, we aim at suggesting the revised model as a general model for specific types of Internet-use disorders, which can then be further specified in future studies with respect to the certain forms of Internet use (e.g., gaming, gambling, pornography, cybersex, social networking, buying/shopping, etc.). Third, we aim at expressing and illustrating the process of the development and maintenance of specific addictive behaviors. By doing this, we explicitly distinguish between predisposing factors, which make individuals vulnerable for developing specific Internet-use disorders, and variables that act as moderators and mediators within the addiction process. Moderator and mediator variables are important components of theoretical models for psychiatric/psychological disorders, because pharmacological and psychological interventions may address moderating and mediating variables effectively, while certain vulnerability factors (e.g., genetic

vulnerability, personality) may be relatively stable (Brand, Laier, et al., 2014). Such theoretical models, or parts there of, may then be transferred into statistical models, which may be tested empirically in future studies. By understanding the mechanisms behind the phenomena, policy, prevention and treatment efforts may be developed and tested on the basis of systematic hypotheses. We aim at suggesting such a theoretical framework for a process model, which hopefully inspires future research and clinical practice.

## **2 Summary of current models on the development and maintenance of Internet-use disorders**

The model for Internet addiction by Brand, Young, et al. (2014) consists of three parts: a model describing the functional/healthy use of the Internet, a model of generalized Internet addiction (see Davis, 2001), and a global model of specific types of Internet addiction. Here, we focus on the revision of the model on specific Internet-use disorders. The specific types refer to an addictive use of one certain genre of applications or sites, such as gaming, gambling, pornography/cybersex, shopping, social networking, or communication. This means we postulate that individuals have a “first-choice use”, which is considered being comparable to the “first-choice drug” in substance-dependent individuals.

This model of specific Internet-use disorders includes psychopathological features (e.g., depression, social anxiety) and dysfunctional personality traits as well as other variables (e.g., stress vulnerability) as factors representing predispositions. Beyond those more global vulnerability factors, we have proposed that persons have specific characteristics, which make them more vulnerable to use certain types of applications or sites addictively. For example, a strong predilection towards gaming or high sexual excitability in general may in part explain why people excessively use

specific applications/sites (i.e., relating to gaming or pornography-viewing, respectively) to experience gratification and pleasure. In terms of a mediation effect, we have also proposed that the predisposing variables may not have a direct impact on the development of a specific Internet-use disorder, but that they are associated with certain Internet-use expectancies and dysfunctional coping styles. Use-expectancies and coping have been considered personal core cognitions and may represent important moderating or mediating variables. As the final part of the model, the use of the first-choice application/site results in experience of gratification and positive reinforcement (Everitt & Robbins, 2016; Piazza & Deroche-Gamonet, 2013). Gratification leads to positive (and partly negative) reinforcement of the dysfunctional coping style, the expectancies about the use of specific Internet applications/sites, and some core characteristics, particularly psychopathological features and the specific preferences. We have argued further that these learning mechanisms may make it increasingly difficult for individuals to exert executive and inhibitory control over their Internet-use behavior.

The theoretical model of Internet-gaming disorder by Dong and Potenza (2014) also includes personal attitudes and cognitive processes. Central to this model is the link between decision-making style in terms of searching for immediate reward despite long-term negative consequences, and motivation-seeking (craving) in terms of a drive to experience pleasure and/or to reduce stress. The third domain involves executive control (inhibition and monitoring) over motivation-seeking, which is hypothesized to be reduced in individuals with Internet-gaming disorder. This assumption is consistent with theories and empirical findings of executive functioning in substance-dependent individuals (Goldstein & Volkow, 2011). In their model, Dong and Potenza (2014) refer to theories on substance addictions, which are reward-centered. One example is the incentive salience theory and the distinction of “liking”

from “wanting” a drug (Berridge, 2007; Berridge, Robinson, & Aldridge, 2009; Robinson & Berridge, 2001, 2008). Dong and Potenza (2014) also included suggestions for treatment interventions, which could target specific cognitive and motivational factors.

Both models, which share several main components, are theoretically plausible, and studies to date have empirically tested parts of them. Previous studies with Internet-gaming disorder and other types of Internet-use disorders could show that certain vulnerability factors, motivation-seeking and craving, cognitive processes and decision-making, are worth considering. On the basis of these two theoretical models and integrating findings from recent studies on Internet-use disorders as well as other research areas, we suggest a revised theoretical process model for specific Internet-use disorders, which is aimed to reflect the addiction process in the development and maintenance of specific Internet-use disorders. This model should be understood as a theoretical framework for Internet-use disorders, although several parts of the model need to be tested empirically in future studies, in particular in investigations comparing different types of Internet-use disorders.

### **3 The Interaction of Person-Affect-Cognition-Execution (I-PACE) model of specific Internet-use disorders**

The I-PACE model includes the following main components: Predisposing variables, affective and cognitive responses to internal or external stimuli, executive and inhibitory control, decision-making behavior resulting in the use of certain Internet applications/sites, and consequences of using the Internet applications/sites of choice. The model is illustrated in figure 1.

((Please insert figure 1 here))

### **3.1 Predisposing variables representing core characteristics of the person: The P-component of the model**

#### **3.1.1 Biopsychological constitution**

Predisposing variables contribute to a person's core characteristics, which may be relatively stable over time. The earliest predisposing factors are genetic factors and other biological determinants of human behavior, such as ontogenetic aspects and early childhood experiences and their resulting biological consequences and effects on learning experiences. With respect to a potential genetic contribution to Internet-use disorders, studies suggest that up to 48% of individual differences in Internet-use-disorder features may be accounted for by genetic factors, although the degrees of heritability estimates vary across investigations (Deryakulu & Ursavas, 2014; M. Li, Chen, Li, & Li, 2014; Vink, Beijsterveldt, Huppertz, Bartels, & Boomsma, 2015). One example for genetic variations that have been linked to Internet-use disorders are related to dopamine systems (in particular polymorphisms *COMT* Val158Met and *ANKK1/DRD2* Taq Ia), as reported by Han et al. (2007). This finding resonates with findings linking candidate polymorphisms to other behavioral addictions, such as pathological gambling (Goudriaan, Oosterlaan, Beurs, & van den Brink, 2004; Potenza, 2013). The serotonin-transporter-linked polymorphic region (5-HTTLPR) of the gene encoding the serotonin transporter (*SLC6A3*) has also been linked to Internet-use disorders (Y. S. Lee et al., 2008). With respect to the cholinergic system as a third potential neurochemical system involved in Internet-use disorders, Montag, Kirsch, Sauer, Markett, and Reuter (2012) reported a link between a genetic variation of the *CHRNA4* gene (linked to the cholinergic nicotine/acetylcholine receptor) and Internet-use-disorder features. However, these studies have typically involved relatively small, incompletely characterized samples and analyses targeted to specific candidate polymorphisms. In summary, although several initial studies give

preliminary evidence for potential genetic contributions to Internet-use disorders, further research is needed (including from genome-wide association studies). It is also likely that individuals with different types of Internet-use disorders represent a heterogeneous group with respect to their genetic profiles. Most studies on genetics have included persons with Internet-gaming disorder or did not differentiate among different types of usage (Weinstein & Lejoyeux, 2015). Future studies should explicitly ask for the “first-choice” use and compare the genetic profiles among different forms of Internet-use disorders (for example, those related to gaming, gambling, pornography viewing, buying, and social networking).

Further biopsychological factors, which may make individuals vulnerable to develop a mental disorder in general or an addictive behavior in particular, are negative early childhood experiences, such as early trauma, emotional or physical abuse, and social isolation. Consistent with this notion, some studies have found correlations between negative early life events and Internet-use disorders (Dalbudak, Evren, Aldemir, & Evren, 2014; Hsieh et al., 2016). Negative life events in early childhood have also been linked to an insecure attachment style, which has also been found related with Internet-use disorders (e.g., Odaci & Çikrikçi, 2014; Schimmenti, Passanisi, Gervasi, Manzella, & Famà, 2014) including problematic Internet pornography use (Kor et al., 2014). One biological correlate of insecure attachment style is lower levels of oxytocin, which have also been associated with developing addictive behaviors (Baskerville & Douglas, 2010; Sarnyai & Kovács, 2014). Consistent with this notion, stressful experiences in early childhood make individuals more vulnerable to react intensively to stress in adolescents and adulthood (Elsley et al., 2015) and to develop mental disorders (Chen & Baram, 2016) and addictive behaviors (Briand & Blendy, 2010). In this context, early childhood experiences in combination with parental styles, familial atmospheres and parents'

own Internet and media use may also have an important impact on children's and adolescents' Internet use and the development of an Internet-use disorder (Lam & Wong, 2015; Zhang, Brook, Leukefeld, & Brook, 2016).

### *3.1.2 Psychopathological features, personality, and social cognitions*

Beyond these vulnerability factors, which develop relatively early or are even prenatally determined, a broad body of literature exists on the correlations and comorbidities of diverse psychopathological features and features of Internet-use disorders. Depression and (social) anxiety disorders as well as attention-deficit/hyperactivity disorder (ADHD) have been considered as three main co-morbid conditions of Internet-use disorders (see meta-analyses by Ho et al., 2014; Prizant-Passal, Shechner, & Aderka, 2016). With respect to personality factors, the most consistent links have been found between Internet-use-disorder features and high impulsivity, low self-esteem, low conscientiousness, high shyness, high neuroticism, a tendency to procrastinate, and low self-directedness (Ebeling-Witte, Frank, & Lester, 2007; Floros, Siomos, Stogiannidou, Giouzevas, & Garyfallos, 2014; Hardie & Tee, 2007; H. K. Kim & Davis, 2009; Koo & Kwon, 2014; K. W. Müller, Beutel, Egloff, & Wöfling, 2014; Niemz, Griffiths, & Banyard, 2005; Sariyska et al., 2014; Thatcher, Wretschko, & Fridjhon, 2008; Wang, Ho, Chan, & Tse, 2015; Weinstein et al., 2015). Social cognitions have been primarily linked to the excessive use of Internet applications/sites which include communication features (for example, social-networking sites and online role-playing games). A perceived lack of social support, feelings of isolation, and loneliness have been considered important in this context (Caplan, 2007; Morahan-Martin & Schumacher, 2003; Odacı & Kalkan, 2010; Pontes, Griffiths, & Patrão, 2014). Again, it is likely that individuals with different types of Internet-use disorders have specific personality profiles. There may be some

commonalities across the different groups. For example, higher ADHD rates and higher impulsivity have been observed in recent meta-analyses (see citations above). However, it is also likely that different types of Internet-use disorders are linked to specific personality traits. One example is the above-mentioned link between social cognitions and using communication applications excessively. Future studies should explicitly address the personality profiles among different forms of Internet-use disorders to explore common and unique correlates of dysfunctional use of certain Internet applications, as it has been done with respect to other fields (e.g., with respect to substance-use disorders).

### *3.1.3 Using motives*

The aforementioned predisposing factors may represent potential risk factors for the development of addictive use of the Internet without considering the specific applications/sites of choice. Although most studies mentioned investigated Internet gaming or did not define the certain use of choice precisely, some disjunctive predispositions may explain individual motives or preferences for using specific applications or sites excessively. Social aspects are particularly relevant for using online communication applications/sites (Kuss & Griffiths, 2011b). Extraversion and openness to experience (Correa, Hinsley, & de Zuniga, 2010) as well as narcissism (T. Ryan & Xenos, 2011) are also considered important in this context. Sexual excitability, on the other hand, should play a more central role in problematic use of Internet pornography and cybersex (Laier & Brand, 2014; Lu, Ma, Lee, Hou, & Liao, 2014). Specific motives may predispose individuals to choose specific forms of Internet use, such as sites for Internet pornography and cybersex (Paul & Shim, 2008; Reid, Garos, & Carpenter, 2011), gaming (Billieux et al., 2013; Demetrovics et al., 2011; King & Delfabbro, 2014; Kuss, Louws, & Wiers, 2012; R. M. Ryan, Rigby, &

Przybylski, 2006; Yee, 2006), or shopping (Kukar-Kinney, Ridgway, & Monroe, 2009). Further subdivisions may also make sense, for example separating the motives for using pornography versus using sex-dating applications or differentiating between shopping sites and sites for online auctions. However, the empirical evidence for such specific predispositions is rare. We argue that certain preferences and motives are relevant for the selection of the first-choice applications/sites. Future research should consider different first-choice applications/sites when investigating using motives in the context of Internet-use disorders.

### ***3.2 Affective and cognitive responses to external or internal stimuli: The A- and C-components of the model***

After reviewing general and specific factors of vulnerability for developing a specific Internet-use disorder, a question remains why some individuals may use certain Internet applications/sites addictively. In other words, what are the mechanisms underlying the decision to use an application/site and which result in a diminished control over Internet use in certain situations?

Situational factors are perceived subjectively, and the subjective perception results in affective and cognitive responses linked to the level of perceived stress (Dickerson & Kemeny, 2004; Koolhaas et al., 2011). Perceived stress resulting from personal conflicts or abnormal mood (e.g., depressive or anxious state, euphoria) may influence cognitive processes, for example, by focusing the attention to short-term rewards and risky decision-making (Starcke & Brand, 2012, in press). Subjective stress responses to situational factors may influence whether or not individuals decide to use the Internet potentially to cope with the associated cognitions and affects (Tavolacci et al., 2013). We propose that both internal and external stimuli may be conditioned within an addiction process (Kalivas & Volkow,

2005; Volkow, Wang, Fowler, & Tomasi, 2012) and may then trigger affective and cognitive processes resulting in the decision to use the Internet application/site of choice. Consistent with this notion, individuals with Internet-gaming disorder may react with changes in mood and other withdrawal symptoms when being confronted with Internet-related cues, and addiction-related cues may be associated with the anticipated gratification or the reduction of withdrawal symptoms (Kaptsis, King, Delfabbro, & Gradisar, 2016; Osborne et al., 2016; Romano, Osborne, Truzoli, & Reed, 2013).

### *3.2.1 Coping*

Experienced stress in daily life and the subsequent use of the Internet as a tool for coping with problematic or stressful life events have also been considered important factors potentially contributing to the development of Internet-use disorders (Tang et al., 2014; Whang, Lee, & Chang, 2003). In particular, tendencies towards impulsive coping strategies when being confronted with daily stress have been considered problematic in this context (Tonioni et al., 2014). Some authors conceptualize Internet-use disorders as dysfunctional coping with everyday life (Kardefelt-Winther, 2014). We propose that individuals who have a greater vulnerability to stress (as predisposing factors) in combination with dysfunctional/impulsive coping strategies may be more inclined to react with an urge for mood regulation when being confronted with a stressful situation. This interaction could then result in a higher probability of using the Internet application/site of choice, if the individual has the (implicit or explicit) expectancy or illusion that using the Internet is stress-relieving or has other Internet-related cognitive biases.

### *3.2.2 Internet-related cognitive biases*

Several cognitive factors, such as general dysfunctional attitudes, are related to features of Internet-use disorders (Noh & Kim, 2016) in combination with Internet-related expectancies or even illusions (i.e., false beliefs about the effects of using certain applications/sites (Taymur et al., 2016)), as well as implicit associations. In the proposed model, these examples of explicit and implicit cognitions about Internet use and their potential effects on individuals are summarized under the term Internet-related cognitive biases. Features of Internet addiction may co-vary positively with both positive expectancies (e.g., to experience pleasure) and avoidance expectancies (e.g., to escape from reality) on a bivariate level (Brand, Laier, et al., 2014; Y. H. Lee, Ko, & Chou, 2014; Turel, Serenko, & Giles, 2011; Xu, Turel, & Yuan, 2012). Moreover, positive metacognitions about Internet use have been shown to mediate the relationship between emotional dysregulation and Internet addiction (Casale, Caplan, & Fioravanti, 2016) as well as the relationship between psychopathological symptoms (depression, social anxiety) and the addictive use of social-networking sites (Wegmann, Stodt, & Brand, 2015). These expectancies have some overlap with motives for using the Internet (see above). A difference is seen in the stability and concreteness of effects. Motives are considered to be relatively stable and predispose the general approach behavior towards certain applications. The concrete expectancies refer to the ideas and thoughts on the concrete effects that using a specific application or site will most likely have in a certain situation. Such expectancies may be explicit or implicit, and one underlying cognitive process may be that using an application frequently and experiencing positive results (e.g., pleasure or escaping from reality) will result in positive (implicit) associations, which may make it more likely to use this application again (reinforcement). Implicit associations have a reliable predictive value in the context of substance addictions (see meta-analysis by Rooke, Hine, & Thorsteinsson, 2008). Such implicit

associations have been demonstrated for Internet gaming (Yen et al., 2011), Internet pornography (Snagowski, Wegmann, Pekal, Laier, & Brand, 2015), and gambling (e.g., Brevers et al., 2013) using a modified version of the Implicit Association Test (Greenwald, McGhee, & Schwartz, 1998). On the basis of these studies on several facets of explicit and implicit cognitions, we propose that Internet-related cognitive biases, which comprise explicit expectancies and illusions and implicit associations, may have an accelerating effect on cue-reactivity and craving, if an individual is confronted with Internet-related cues and other situational variables (e.g., negative or very positive moods, stress).

### *3.2.3 Cue-reactivity and craving*

One main process behind the diminished behavioral control is craving, which was also defined as motivation-seeking in the model by Dong and Potenza (2014). Craving originally referred to a difficult-to-resist urge to consume a substance. Craving can be triggered by cue-reactivity, which is the result of the confrontation with conditioned addiction-related stimuli (Breiner, Stritzke, & Lang, 1999; Carter & Tiffany, 1999). Cue-reactivity is developed on the basis of (associative) learning mechanisms, in particular conditioning processes (Carter & Tiffany, 1999; Loeber & Duka, 2009; Tiffany, Carter, & Singleton, 2000), which provide the main physiological, emotional, and motivational basis for craving (Robinson & Berridge, 1993, 2000). The concepts of cue-reactivity and craving have been transferred from research on substance addictions to that on behavioral addictions, for example with respect to gambling disorder (e.g., Potenza, 2008; Potenza et al., 2003; Wöfling et al., 2011). Several fMRI studies have investigated brain correlates of cue-reactivity and craving in individuals with gambling disorder (Crockford, Goodyear, Edwards, Qickfall, & el-Guebaly, 2005; Goudriaan, de Ruiter, van den Brink, Oosterlaan, &

Veltman, 2010; Kober et al., 2016; Miedl, Büchel, & Peters, 2014; Potenza et al., 2003; Wulfert, Maxson, & Jardin, 2009). These studies typically observe an involvement of the ventral striatum (and partially further structures of the expanded limbic system) in the experience of craving when being confronted with addiction-related cues. More recently, neural correlates of cue-reactivity and craving, also consistently focusing on the ventral striatum, were demonstrated in subjects with Internet-gaming disorder (Ahn, Chung, & Kim, 2015; Ko et al., 2009; Liu et al., 2016; Thalemann, Wöfling, & Grüsser, 2007), hypersexual behavior (Klucken, Wehrum-Osinsky, Schweckendiek, Kruse, & Stark, 2016; Voon et al., 2014), and Internet-pornography-use problems (Brand, Snagowski, Laier, & Maderwald, 2016). The findings fit well with previous behavioral investigations of the important role of craving and the anticipation of sexual gratification in individuals with cybersex problems (Brand et al., 2011; Laier, Pawlikowski, Pekal, Schulte, & Brand, 2013), and demonstrate the involvement of the ventral striatum in the process of cue-reactivity and craving in behavioral addictions.

#### *3.2.4 Urge for mood regulation*

When being confronted with abnormal mood, withdrawal symptoms, or craving, an urge to regulate the experienced mood may develop. The process of emotion regulation is an important consideration across multiple psychopathological conditions including addictions (Aldao, Nolen-Hoeksema, & Schweizer, 2010; Gross & Jazaieri, 2014; Thorberg & Lyvers, 2006). It has been reported that addictive behaviors may be used in a dysfunctional fashion to cope with experienced aversive affective responses to internal or external cues; e.g., with respect to smoking, alcohol drinking, and use of Internet pornography and performance of online gaming or social networking (Holahan, Moos, Holahan, Cronkite, & Randall, 2001; Hormes, Kearns, &

Timko, 2014; Kuss, 2013; Laier & Brand, 2014; J.-R. Li, Wang, & Lin, 2012; Shapiro, Jamner, Davydov, & James, 2002). Abstinent individuals in recovery from drug addictions may be at elevated risk for relapse in situations in which they are confronted with internal or external cues associated with former drug intake (Welberg, 2013). We propose that the urge for mood regulation is an important factor within the development of Internet-use disorders because it may influence the decision to use certain Internet applications/sites in the early stages of an addiction process. Furthermore, a role may become more important later within the addictive process since perception of experienced problems should lead to greater aversive moods, while coping skills decline in favor of dysfunctional coping by using the Internet applications/sites of choice.

### *3.2.5 Attentional biases*

Attentional biases and their relationships with craving responses have been studied in substance addictions (e.g., Christiansen, Schoenmakers, & Field, 2015; Field & Cox, 2008; Field, Munafò, & Franken, 2009). Implicit cognitions, in particular approach and avoidance tendencies, have been related to craving responses in substance-dependent individuals (e.g., R. W. Wiers & Stacy, 2006). The idea that attentional biases may guide addictive behaviors fits well with recent dual-mode theories of addictive behaviors (e.g., Bechara, 2005; Evans & Coventry, 2006; Stacy & Wiers, 2010). These approaches share main views on the nature of addiction, which is that addictive behaviors may result from the interaction of two types of processes. The first type is an impulsive or relatively automatic processing mode, and the second type is a relatively controlled and reflective processing mode. This general approach of viewing addictive behaviors as the outcome of an impulsive and a deliberate cognitive processing mode is consistent with current decision-making

theories (e.g., Schiebener & Brand, 2015) and cognitive psychology models of dual-processing in reasoning and thinking (Evans, 2003; Kahneman, 2003; Stanovich & West, 2000).

Evidence for attentional bias has been observed in problem-gamblers (Ciccarelli, Nigro, Griffiths, Cosenza, & D'Olimpio, 2016). Attentional bias in individuals with Internet-gaming problems was demonstrated recently (Jeromin, Nyenhuis, & Barke, 2016), with attentional bias measured using two instruments which have been widely used in studies of substance addictions: the Addiction Stroop Task and visual-probe tests (Field & Cox, 2008). Individuals with Internet-gaming problems as compared to those without reacted more slowly to computer-related words as compared to neutral words during addiction-Stroop performance, which can be regarded as an attentional bias towards the addiction-related stimuli. The results fit with those from studies of Internet use and video-game playing, which also used an Addiction Stroop Task (Metcalf & Pammer, 2011; van Holst et al., 2012), although in the study by van Holst et al. (2012) reaction times for addiction-related cues and neutral words were not different. With respect to the visual-probe findings, no differences in reaction times were found in either studies (Jeromin et al., 2016; van Holst et al., 2012), but participants made more errors for targets in the condition with computer-related words, indicating a potential interference between attentional bias and correctly identifying the target position. Even clearer findings were observed in patients with hypersexual behavior compared to healthy volunteers during performance of a visual-probe task; individuals with hypersexual behaviors showed greater attentional bias to explicit sexual stimuli relative to neutral pictures (Mechelmans et al., 2014).

Another line of research within the field of affective-attentional processes in addicted individuals is the relationship between craving and the tendency to

approach or avoid addiction-related stimuli (Breiner et al., 1999). Studies suggest a multi-dimensional model for alcohol addiction that focuses on an evaluative space in the situation when being confronted with addiction-related stimuli. Positive or negative expectancies towards the effects of drug intake may influence the tendency to approach or to avoid drug-related cues. Positive expectancies should result in approach tendencies while negative expectancies should result in avoidance tendencies. The approach/avoidance framework is also in line with the aforementioned dual-process models of addictive behaviors. One task that has been frequently used in alcohol-use research to measure approach and avoidance tendencies is the Approach-Avoidance Task, which was originally developed by Rinck and Becker (2007) to investigate individuals with anxiety disorder (spider phobia). The task includes a physical movement via joystick and participants have to pull stimuli presented on a computer screen towards themselves (approach condition) or push them away (avoidance condition) from themselves as fast as possible. Multiple studies indicate that addicted subjects react faster when they have to approach the drug-related stimuli as compared to non-addicted individuals or compared to the avoidance condition (Cousijn et al., 2012; Cousijn, Goudriaan, & Wiers, 2011; C. E. Wiers et al., 2013). Using the Approach-Avoidance Task, Snagowski and Brand (2015) found that individuals with problematic Internet pornography use (in an analogue sample) can be linked to both approach and avoidance tendencies, as they found a quadratic relationship in their sample of pornography users. Although these results should be considered cautiously, since they need to be replicated and transferred to other types of Internet-use disorders, it seems worth considering such approach and avoidance tendencies as potential mechanisms underlying addictive use of certain Internet applications/sites.

In summary, predisposing factors together with dysfunctional coping styles, Internet-use expectancies, illusions, and implicit associations may influence the intensities of cue-reactivity and craving and other specific cognitive and affective processes, such as attentional biases and approach tendencies towards addiction-related stimuli. Consistent with some of the findings on potential interactions, although studies addressing explicitly interaction effects between variables are still scarce, we propose that the predisposing variables act in concert with coping styles and Internet-related biases resulting in specific patterns of affective and cognitive responses in specific situations. The affective and cognitive responses, as results of interaction effects, include cue-reactivities, cravings, urges for mood regulation, and attentional biases. We consider these as being important processes that impact decisions to use certain applications/sites. However, we also propose that mediating variables may exist between affective and cognitive responses and the decision to use the Internet, and these mediating factors may reside in the domains of inhibitory control and executive functioning.

### ***3.3 Executive functions, inhibitory control, and the decision to use certain applications/sites: The E-component of the model***

The potential impact of reduced executive functioning and reduced inhibitory control were central ingredients of the model on Internet-gaming disorder by Dong and Potenza (2014) and also of the model by Brand, Young, et al. (2014), although this was not included explicitly in the figure, but described in the text (Brand, Young, et al., 2014). The idea that executive functions contribute importantly to the development and maintenance of specific Internet-use disorders is based on neuropsychological and neuroscientific research and theories of substance addictions (Bechara, 2005; Goldstein et al., 2009; Goldstein & Volkow, 2002, 2011;

Kalivas & Volkow, 2005; Koob & Volkow, 2010; Volkow & Fowler, 2000; Volkow, Fowler, Wang, & Goldstein, 2002; Volkow et al., 2012). These models propose that reduced function of the prefrontal cortex is linked to impaired response inhibition and salience attribution (IRISA-model) in individuals with addictions. A main characteristic of this model is the increased salience to drug-related stimuli and – simultaneously – decreased sensitivity to natural, non-substance-related reinforcers. As a consequence of this interaction, diminished control over the addictive behavior and reduced inhibition of disadvantageous decision-making occur (cf. Goldstein & Volkow, 2011). We argue that the diminished control over decision-making in the context of addictions can be transferred to behavioral addictions and specific Internet-use disorders.

Executive functions, inhibitory control, and decision-making have been studied in the context of Internet-use disorders, with a focus on Internet-gaming disorder in particular (e.g., Dong, Hu, Lin, & Lu, 2013; Pawlikowski & Brand, 2011; Sun et al., 2009). Findings regarding inhibitory control in individuals with Internet-use disorders are mixed, although the majority of studies found at least mild executive reductions in individuals with Internet-use disorders (Dong, Hu, et al., 2013; Dong, Lu, Zhou, & Zhao, 2010; Dong, Zhou, & Zhao, 2011; Sun et al., 2009; van Holst et al., 2012). The same appears true for decision-making, as some studies have found no general differences between subjects with and without Internet-use disorders during decision-making under ambiguous conditions, as measured with the Iowa Gambling Task (Yao et al., 2015), while others have found affected individuals performing inferiorly to healthy volunteers (Sun et al., 2009). More consistently, significant decision-making reductions were found in tasks assessing decisions under risk conditions (Dong & Potenza, 2016; Pawlikowski & Brand, 2011; Seok, Lee, Sohn, & Sohn, 2015; Yao et al., 2015). When comparing individuals with either Internet-use or alcohol-use

disorders, both groups had comparable levels of performance in executive-function tasks, and both groups scored significantly lower in comparison to healthy volunteers (Z. Zhou, Zhu, Li, & Wang, 2014).

Most studies to date of inhibitory control using the Go/No-Go Task have employed versions with neutral stimuli (i.e., without addiction-related stimuli) and observed no reductions in behavioral performance (Ding et al., 2014), although results are mixed across the existing studies (see meta-analysis by Smith, Mattick, Jamadar, & Iredale, 2014). As with studies on attentional biases, studies may be more informative and findings may be more consistent if addiction-related stimuli were used. We hypothesize that subjects with specific Internet-use disorders may have difficulties in inhibiting responses to stimuli, which represent their first-choice-use, as has been shown in binge drinkers (Czapla et al., 2015) and substance-dependent individuals (e.g., Pike, Stoops, Fillmore, & Rush, 2013). In this context, Z. Zhou, Yuan, and Yao (2012) used a shifting task with cues representative for Internet games and found reductions in response inhibition and mental flexibility. In a cue-specific version of the Go/No-Go Task, cue-related reductions of inhibitory control in individuals with Internet-gaming disorder were reported (Yao et al., 2015). Another example is the study by Nie, Zhang, Chen, and Li (2016) showing impaired response inhibition and working memory in adolescents with Internet-use disorder in the Stop Signal Task and 2-back Task including Internet-related words as cues. Consistent with this finding, Laier, Pawlikowski, and Brand (2014) used a modified Iowa Gambling Task with pornographic and neutral pictures on the advantageous and disadvantageous card decks (and *vice versa* in the other group of subjects). In a sample of male pornography users, those individuals who performed the task with pornographic pictures on the disadvantageous card decks continued choosing cards from these decks despite receiving high losses. This effect was accelerated in

participants who reported high subjective craving after the presentation of pornographic pictures in an additional experimental task.

The findings on reduced executive functioning and inhibitory control, perhaps as a consequence of cue-reactivity and craving, are consistent with results obtained from neuroimaging studies (cf. Kuss & Griffiths, 2012; Meng, Deng, Wang, Guo, & Li, 2015; Sepede et al., 2016). Structural differences in individuals with and without Internet-gaming disorder have been reported in both gray and white matter in prefrontal brain areas and additional brain regions, such as limbic structures (e.g., Hong, Kim, et al., 2013; Hong, Zalesky, et al., 2013; H. Wang et al., 2015; Y. Zhou et al., 2011). Functional brain correlates of Internet-gaming disorder are also reported in the prefrontal cortex and limbic structures (Dong, Devito, Du, & Cui, 2012; Dong, Hu, et al., 2013; Dong, Lin, Zhou, & Lu, 2014). Changes in dopaminergic systems have also been proposed (S. H. Kim et al., 2011), which may relate to reinforcement processing (Jović & Đinđić, 2011). Studies are also beginning to bring together findings from neuropsychological investigations and neuroimaging assessments in individuals with Internet-gaming disorder or problematic gaming behavior, which show that deficits in executive functions and inhibitory control are related to functional changes in fronto-striatal circuits (Luijten, Meerkerk, Franken, van de Wetering, & Schoenmakers, 2015; Seok et al., 2015; Yuan et al., 2016).

Taken together, reductions in executive functions, inhibitory control, and decision-making are present in individuals with Internet-use disorders or in individuals who appear at elevated risk for developing addictive patterns of Internet use, particularly in situations when they are confronted with Internet-addiction-related cues. The neural correlates of Internet-gaming disorder and other Internet-use disorders (e.g., Brand et al., 2016) may reflect a maladaptive interaction of cue-reactivity/craving and reduced prefrontal/executive functioning, as suggested for

substance addictions (Goldstein & Volkow, 2011; Koob & Volkow, 2010; Volkow & Fowler, 2000; Volkow et al., 2002). We propose that a dysfunctional interaction between poor executive control and situationally accelerated reward-seeking, as a result of cue-reactivity and craving, may promote disadvantageous decision-making. The decision to use certain Internet applications/sites to reduce craving and to increase mood is regarded as being characterized by seeking a short-term attractive behavior that results in the experience of gratification despite negative long-term consequences. This hypothesized type of dysfunctional interaction between executive control and reward-seeking has been emphasized recently by the fMRI study of Dong, Lin, Hu, Xie, and Du (2015). They used resting-state fMRI and showed decreased functional connectivity in the so-called executive-control network (including lateral prefrontal and parietal regions) in individuals with Internet-gaming disorder compared to healthy volunteers. In addition, the individuals with Internet-gaming disorder showed increased functional connectivity in reward-associated networks (including ventral striatum and orbitofrontal cortex). Dong and colleagues propose that the imbalance between executive control and reward networks represent a mechanism seen in individuals with Internet-gaming disorder, with reductions in executive control leading to a diminished inhibition of motivation-seeking and craving, resulting in excessive Internet gaming. We agree with this interpretation, as reflected in our model in the path from affective and cognitive responses over reductions in executive function and inhibitory control to disadvantageous decision-making. Future studies may investigate decision-making, executive functions, and inhibitory control with and without addiction-related stimuli and compare performances across different types of Internet-use disorders. Such studies could provide a more complete picture of how specific cognitive processes

may be involved in the development and maintenance of specific Internet-use disorders.

### ***3.4 Consequences resulting from using the Internet applications/sites of choice***

The decision to use certain applications/sites and the behavior of using them may lead to short-term positive experiences and gratification, at least in the early stages of the addiction process. In addition, and perhaps even more importantly, the use of certain Internet applications/sites and the gratification received should also lead to an increase in cue-reactivity and craving as responses to certain stimuli, as a result of both Pavlovian- and instrumental-conditioning processes. The importance of conditioning in the development of addictive behaviors has been suggested theoretically, for example within the incentive sensitization theory (Berridge et al., 2009; Robinson & Berridge, 1993, 2001, 2008), and has been demonstrated empirically in substance addictions (Duka et al., 2011; Hogarth, Dickinson, & Duka, 2010; Hogarth, Dickinson, Hutton, Elbers, & Duka, 2006; Loeber & Duka, 2009), for example using the Pavlovian Instrumental Transfer Task (Hogarth, Dickinson, Wright, Kouvaraki, & Duka, 2007). Recent data suggest that similar conditioning processes are also involved in developing cue-reactivity and craving in the context of Internet-pornography-use disorder (Klucken et al., 2016; Snagowski, Laier, Duka, & Brand, in press). Although for other types of Internet-use disorders empirical evidence for conditioning processes is still missing, we propose that experiencing gratification due to using the Internet applications/sites of choice leads to positive reinforcement, which is the basis for developing stabilizing cue-reactivity and craving. Consistent with this notion, we also propose that based on reinforcement learning, the dysfunctional coping styles and the Internet-related cognitive biases are positively and partly negatively reinforced and therefore strengthened. All of these

reinforcement mechanisms may make it more likely that individuals use the applications/sites of choice repeatedly. The mechanisms may also make it more likely that the applications/sites of choice are used in many situations, akin to what is seen in substance addictions. Due to conditioning processes in substance addictions, a generalization of situational features triggering cue-reactivity and craving is developed and the addictive behavior becomes habitual and/or compulsive (cf. Everitt, 2014; Everitt & Robbins, 2005, 2016). The suggested reinforcement cycle, which represents the temporal dynamic within the middle/grey part of the model (figure 1), is presented in figure 2.

((please insert figure 2 about here))

The addiction process in general has been proposed to involve transitions from more voluntary and impulsive drug consumption to a more habitual or compulsive pattern of use, and that within this process, the positive and recreational feelings linked to drug intake may become less important compared to experiencing the direct drug effects (Everitt & Robbins, 2016; Piazza & Deroche-Gamonet, 2013). We propose that in the early stages of the process of specific Internet-use disorders, gratification is a main, but not exclusive, driving force that leads to changes in affective and cognitive responses to Internet-addiction-related stimuli. As the addiction process progresses, the level of experienced gratification decreases. Concurrently, the level of compensating effects increases in the course of the addiction. As control over the use of specific Internet applications/sites diminishes, there may be an increase in negative consequences, which may include social isolation and loneliness, conflicts with parents or peers, feelings of being misunderstood, feelings of emptiness, and other negative emotions and experiences.

These feelings and losses of social contacts or other problems may be further exacerbated by repeatedly using the Internet applications/sites of choice, with gratification becoming less important, and compensation becoming more important. The hypothesized shift from gratification to compensation in the addiction process is summarized in figure 3.

((please insert figure 3 about here))

#### **4 Clinical implications**

Given the recognized clinical significance of Internet-use disorders, clinicians and researchers have been developing specific treatments for individuals with Internet-use problems (Young, 2009), although only Internet-gaming disorder has been included in 2013 as a research diagnosis in section III in the DSM-5. Both pharmacological and psychological treatments have been suggested, similar to what is recommended for gambling disorder and other behavioral addictions (e.g., Grant, Schreiber, & Odlaug, 2013; Yau & Potenza, 2015), and initial studies suggest varying degrees of efficacy (Cash et al., 2012; Santos et al., 2016; Winkler, Dörsing, Rief, Shen, & Glombiewski, 2013; Young, 2013). Cognitive-behavioral therapy for Internet addiction (CBT-IA) was introduced by Young (2011), which is reported currently as the method of choice (Cash et al., 2012; Winkler et al., 2013). However, large-scale randomized controlled trials are needed to evaluate further the efficacies of interventions and additional studies are needed to determine their translational feasibilities into non-research settings. Additionally, given that no medication has an indication for Internet-use disorders, additional research is needed in pharmacotherapy development.

The theoretical framework proposed may be used to promote future clinical interventions. Given that some predisposing factors may not be malleable (e.g., genetics, early childhood experiences) and others may be difficult to change (e.g., psychopathological vulnerability factors, personality), we propose that treatments should primarily address moderating and mediating variables, which may be theoretically modified by pharmacotherapy or psychotherapy. In this context, it is also important to note that even some of the predispositions, such as genetic constitution and stress vulnerability, interact with other moderating and mediating variables. For example, stress vulnerability may moderate the relationship between executive functions and decision-making (Starcke & Brand, in press), and may therefore have an impact on therapy success. Those predisposing factors should be examined in treatment contexts, to better observe potential interactions between personal factors within the treatment process. Variables that may be addressed directly in CBT include coping styles, Internet-related expectancies, attentional biases, cue-reactivities and cravings, as well as executive functions and inhibitory control.

In CBT-IA, an individual's Internet behavior is analyzed and monitored with respect to its situational, emotional, and cognitive contexts. In addition, the subsequent reinforcing effects of Internet use are considered. This process helps to generate understanding of the cognitive assumptions and distortions relating to Internet use and situational triggers. This first phase of CBT-IA considers several variables included in the theoretical model, in particular coping with situations in daily life that are high-risk situations for using the Internet excessively, expectancies and illusions about Internet use and reinforcing effects of Internet use. Afterwards, using methods of cognitive restructuring and reframing, the Internet-related cognitive biases may be targeted.

Given that both explicit and implicit cognitions, as well as an individual's conditionability, may interact with each other (Bernardin, Maheut-Bosser, & Paille, 2014; Forrest, Smith, Fussner, Dodd, & Clerkin, 2016; R. W. Wiers, Boelema, Nikolaou, & Gladwin, 2015), not only the explicit (verbalized) expectancies but also implicit cognitions should be addressed by treatments. S. Lee and Lee (2015) suggested that basic tenets of implicit and explicit cognitions, consistent with roles of approach/avoidance tendencies, could be implemented into therapy as a part of a patient's psychoeducation. Studies of substance addictions suggest that dysfunctional effects of implicit cognitions can potentially be retrained, for example, to increase the probability that experiencing craving might result in avoidance rather than approach tendencies (Eberl et al., 2013a, 2013b; R. W. Wiers, Eberl, Rinck, Becker, & Lindenmeyer, 2011). One way to transfer the concept of retraining to the treatment of Internet-use disorders could be to adapt existing training programs in which patients learn to avoid Internet-related stimuli (e.g., by pushing them away with a joystick, as this is a common training method). However, it should be noted that systematic studies would need to be conducted in order to identify the optimal number of training sessions (Eberl et al., 2013b), and also to evaluate their efficacy. Further methods could consider implicit associations, as has been done for alcohol-use disorders (Houben, Schoenmakers, & Wiers, 2010; C. E. Wiers et al., 2015). However, evidence for the effectiveness of such methods is limited.

Attentional biases may also be decreased in attentional retraining programs (e.g., Christiansen et al., 2015; Schoenmakers et al., 2010). Closely related to this, it was suggested that an individual's ability to inhibit specific actions can be modified through training (e.g., Bowley et al., 2013; Houben & Jansen, 2011; Houben, Nederkoorn, Wiers, & Jansen, 2011), for example, by using modified versions of the Go/No-Go Task. An adoption of these techniques may be beneficial for increasing

inhibitory control and executive functioning and may be included into the treatment of Internet-use disorders, if future studies demonstrated them leading to treatment success. Conditioning processes, which represent main processes underlying Internet-use disorders, may be addressed by methods of cue-exposure therapy (Park et al., 2015). While cue-exposure therapy may not extinguish existing associations, the intensity of experienced craving may be reduced (Pericot-Valverde, García-Rodríguez, Gutiérrez-Maldonado, & Secades-Villa, 2015), which is consistent with current neuroimaging findings on the reduction of cue-reactivity due to cue-exposure therapy in abstinent alcohol-dependent individuals (Vollstädt-Klein et al., 2011), although its effectiveness is discussed controversially (Everitt & Robbins, 2016).

In summary, we propose that it is important to consider assessing individuals' cognitive functions, including attentional biases, implicit and explicit cognitions, executive functions and inhibitory control capacities, in the context of clinical treatment. We also propose that including neuropsychological training with a focus on Internet-specific control processes may increase the likelihood of positive outcomes relating to CBT in the context of Internet-use disorders.

## **5 Critical comments and future directions**

Although the field of research on Internet-use disorders has grown rapidly over the last two decades and many studies on the phenomena exist, there still exist sizable gaps of knowledge, particularly with respect to treatment interventions. Multiple aspects of existing studies limit our current knowledge. First, most empirical studies concentrate on Internet-gaming disorder or do not differentiate among different types of Internet usage. Second, many previous studies have addressed single variables, such as personality or genetic correlates and cognitive functions, relatively isolated from each other and for one form of Internet-use disorders only. Third, most studies

have a cross-sectional design, limiting insight into the development and maintenance of Internet-use disorders. There are some longitudinal studies (e.g., Strittmatter et al., 2016; Zhang et al., 2016), but these are few and limited (e.g., with respect to the time of assessment). Fourth, most studies concentrate on adolescents and young adults and do not include questions regarding early development of the disorders, such as parental and family features. Fifth, gender aspects have not been addressed systematically in meta-analyses because most studies concentrating on Internet-gaming disorder (and also those focusing on Internet-pornography viewing) include mainly or solely male participants.

Given this lack of systematic research, the model suggested cannot be viewed as final. Although we have tried to include results from current research from different areas, not all aspects included in the model are empirically tested for all types of Internet-use disorders. In addition, results are mixed for some aspects, for example personality or decision-making, as we have discussed in the respective sections. However, we believe that the model suggested has the potential to influence future research by providing an explicitly framework for testing of hypotheses with respect to interactions of specific features including personal characteristics and cognitive and affective processes.

In future studies, interactions between core personal, and cognitive and affective features should be considered more systematically. In more detail, a better understanding of the interactions of personality and other trait variables, and cognitive and affective variables, which may develop within an addiction process, such as cue-reactivity, craving, attentional bias, and executive functions, is needed. Investigating the interactions of these variables instead of studying these variables separately seems to be very important in order to contribute to a better understanding of the nature and dynamics of Internet-use disorders. Although Internet-gaming

disorder is the type of Internet-use disorder that is arguably most prominent in clinical practice and the published research literature, it is also important to consider other potential types of Internet-use disorders and compare profiles and underlying mechanism among the different types. For example, Internet-gaming, Internet-gambling, Internet-pornography-use behaviors and disorders, amongst others, warrant consideration and attention. The lack of knowledge in these areas may have been a limiting factor with respect to considering Internet-use disorders in DSM-5, and may hinder efforts relating to how Internet-use disorders are considered in other classification systems like ICD-11.

From the current state of research, we suggest to include Internet-use disorders in the upcoming ICD-11. It is important to note that beyond Internet-gaming disorder, other types of applications are also used problematically. One approach could involve the introduction of a general term of Internet-use disorder, which could then be specified considering the first-choice application that is used (for example Internet-gaming disorder, Internet-gambling disorder, Internet-pornography-use disorder, Internet-communication disorder and Internet-shopping disorder). The general term Internet-use disorder may also cover mixed forms of problematic or addictive use of more than one application (for example, a mixed type of Internet-gaming and Internet-gambling disorder). Based on the data we present in this synthetic review, we hypothesize that, although the evidence is still inconsistent in detail and future studies are needed, different types of Internet-use disorders likely share some core aspects and the I-PACE model incorporates these similarities in a structured framework for direct and systematic examination.

## 6 Conclusion

The Interaction of Person-Affect-Cognition-Execution (I-PACE) model for specific Internet-use disorders is aimed at providing a theoretical framework, which distinguishes between predisposing factors and moderating and mediating variables. Coping styles and Internet-related cognitive biases are mainly conceptualized as moderating variables that may influence associations between predisposing factors and aspects of Internet-use disorders. Coping styles and cognitive biases may also act as mediating variables, which are influenced by, for example, psychopathologies and personality/temperamental characteristics. We further hypothesize the existence of moderated mediation effects between predisposing factors and the moderators/mediators coping styles and Internet-related cognitive biases. Affective and cognitive responses (e.g., cue-reactivities and cravings, attentional biases) to certain situational stimuli are referred to as mediating variables. These reactions should be influenced by the predispositions, but even more strongly by coping styles and Internet-related cognitive biases, and they are considered to develop within the addiction process as a result of conditioning processes in terms of positive and negative reinforcement. These affective and cognitive responses to situational stimuli may reduce inhibitory control and executive functioning, which then contribute to the decision to use the Internet applications/sites of choice. This process is hypothesized as a partial mediation, meaning that also direct effects from affective and cognitive responses to decisions to use certain applications/sites are strong by themselves, but that these effects are partially mediated by reductions in inhibitory control as a result of the responses to the situational features. In summary, the suggested I-PACE model is aimed at summarizing the mechanisms underlying the development and maintenance of specific Internet-use disorders in terms of a process model indicating the temporal dynamics of the addiction process. The ventral striatum and prefrontal

brain areas are considered important neural contributors to the interaction of cue-reactivity and craving with reduced executive functions and diminished decision-making skills in individuals with specific Internet-use disorders. Although the components and processes within the I-PACE model are derived from previous theoretical and empirical studies, the hypothesized mechanisms should be investigated systematically in future studies. The model's assumptions should be specified in more detail for the specific types of Internet-use disorders, for example Internet-gaming, Internet-gambling, Internet-pornography-use, Internet-shopping, and Internet-communication disorders. We hope that the I-PACE model of specific Internet-use disorders inspires future research and clinical practice and is helpful for formulating clear research hypotheses within a rapidly developing and important scientific field.

### **Funding**

Dr. Potenza received support from the National Center for Responsible Gaming and the National Center on Addiction and Substance Abuse. The contents of the manuscript are solely the responsibility of the authors and do not necessarily represent the official view of any of the funding agencies.

### **Declarations of Interests**

The authors report that they have no financial conflicts of interest with respect to the content of this manuscript. Dr. Potenza has received financial support or compensation for the following: Dr. Potenza has consulted for and advised Boehringer Ingelheim, Ironwood, Lundbeck, INSYS, Shire, RiverMend Health, Opiant/Lakelight Therapeutics and Jazz Pharmaceuticals; has received research

support from the NIH, Veteran's Administration, Mohegan Sun Casino, the National Center for Responsible Gaming, and Pfizer, Forest Laboratories, Ortho-McNeil, Psyadon, Oy-Control/Biotie and Glaxo-SmithKline pharmaceuticals; has participated in surveys, mailings or telephone consultations related to drug addiction, impulse control disorders or other health topics; has consulted for law offices and the federal public defender's office in issues related to impulse control disorders; provides clinical care in the Connecticut Department of Mental Health and Addiction Services Problem Gambling Services Program; has performed grant reviews for the NIH and other agencies; has edited journals and journal sections; has given academic lectures in grand rounds, CME events and other clinical or scientific venues; and has generated books or book chapters for publishers of mental health texts. The other authors report no biomedical financial interests or other conflicts of interest.

## References

- Ahn, H. M., Chung, H. J., & Kim, S. H. (2015). Altered brain reactivity to game cues after gaming experience. *Cyberpsychology, Behavior and Social Networking*, *18*, 474-479. doi:10.1089/cyber.2015.0185
- Aldao, A., Nolen-Hoeksema, S., & Schweizer, S. (2010). Emotion-regulation strategies across psychopathology: A meta-analytic review. *Clinical Psychology Review*, *30*, 217-237. doi:10.1016/j.cpr.2009.11.004
- APA. (2013). *Diagnostic and Statistical Manual of Mental Disorders 5th Edition*. Washington DC: APA.
- Baskerville, T. A., & Douglas, A. J. (2010). Dopamine and oxytocin interactions underlying behaviors: Potential contributions to behavioral disorders. *CNS Neuroscience and Therapeutics*, *16*, 92–123.
- Bechara, A. (2005). Decision making, impulse control and loss of willpower to resist drugs: A neurocognitive perspective. *Nature Neuroscience*, *8*, 1458-1463. doi:10.1038/nn1584
- Bernardin, F., Maheut-Bosser, A., & Paille, F. (2014). Cognitive impairments in alcohol-dependent subjects. *Frontiers in Psychiatry*, *5*, 1-6. doi:10.3389/fpsy.2014.00078
- Berridge, K. C. (2007). The debate over dopamine's role in reward: The case for incentive salience. *Psychopharmacology*, *191*, 391-431.
- Berridge, K. C., Robinson, T. E., & Aldridge, J. W. (2009). Dissecting components of reward: 'Liking', 'wanting', and learning. *Current Opinions in Pharmacology*, *9*, 65-73. doi:10.1016/j.coph.2008.12.014
- Billieux, J., Van Der Linden, M., Achab, S., Khazaal, Y., Paraskevopoulos, L., Zullino, D., & Thorens, G. (2013). Why do you play World of Warcraft? An in-depth exploration of self-reported motivations to play online and in-game behaviours in the virtual world of Azeroth. *Computers in Human Behavior*, *29*, 103–109.
- Bowley, C., Faricy, C., Hegarty, B., Johnston, S., Smith, J. L., Kelly, P. J., & Rushby, J. A. (2013). The effects of inhibitory control training on alcohol consumption, implicit alcohol-related cognitions and brain electrical activity. *International Journal of Psychophysiology*, *89*, 342-348. doi:10.1016/j.ijpsycho.2013.04.011
- Brand, M., Laier, C., Pawlikowski, M., Schächtle, U., Schöler, T., & Altstötter-Gleich, C. (2011). Watching pornographic pictures on the Internet: Role of sexual arousal ratings and psychological-psychiatric symptoms for using Internet sex sites excessively. *Cyberpsychology, Behavior, and Social Networking*, *14*, 371-377. doi:10.1089/cyber.2010.0222
- Brand, M., Laier, C., & Young, K. S. (2014). Internet addiction: Coping styles, expectancies, and treatment implications. *Frontiers in Psychology*, *5*, 1256. doi:10.3389/fpsyg.2014.01256
- Brand, M., Snagowski, J., Laier, C., & Maderwald, S. (2016). Ventral striatum activity when watching preferred pornographic pictures is correlated with symptoms of Internet pornography addiction. *Neuroimage*, *129*, 224-232.
- Brand, M., Young, K. S., & Laier, C. (2014). Prefrontal control and Internet addiction: A theoretical model and review of neuropsychological and neuroimaging findings. *Frontiers in Human Neuroscience*, *8*, 375. doi:10.3389/fnhum.2014.00375
- Breiner, M. J., Stritzke, W. G. K., & Lang, A. R. (1999). Approaching avoidance. A step essential to the understanding of craving. *Alcohol Research & Therapy*, *23*, 197-206.

- Brevers, D., Cleeremans, A., Hermant, C., Tibboel, H., Kornreich, C., Verbanck, P., & Noël, X. (2013). Implicit gambling attitudes in problem gamblers: Positive but not negative implicit associations. *Journal of Behavioral Therapy and Experimental Psychiatry*, *44*, 94-97. doi:10.1016/j.jbtep.2012.07.008
- Briand, L. A., & Blendy, J. A. (2010). Molecular and genetic substrates linking stress and addiction. *Brain Research*, *1314*, 219–234.
- Caplan, S. E. (2007). Relations among loneliness, social anxiety, and problematic Internet use. *Cyberpsychology & Behavior*, *10*, 234-242. doi:10.1089/cpb.2006.9963
- Carter, B. L., & Tiffany, S. T. (1999). Meta-analysis of cue-reactivity in addiction research. *Addiction*, *94*, 327-340.
- Casale, S., Caplan, S. E., & Fioravanti, G. (2016). Positive metacognitions about Internet use: The mediating role in the relationship between emotional dysregulation and problematic use. *Addictive Behaviors*, *59*, 84–88. doi:10.1016/j.addbeh.2016.03.014
- Cash, H., Rae, C. D., Steel, A. H., & Winkler, A. (2012). Internet addiction: A brief summary of research and practice. *Current Psychiatry Reviews*, *8*, 292-298. doi:10.2174/157340012803520513
- Chamberlain, S. R., Lochner, C., Stein, D. J., Goudriaan, A. E., van Holst, R. J., Zohar, J., & Grant, J. E. (2015). Behavioural addiction - A rising tide? *European Neuropsychopharmacology*, *S0924-977*, 266-267. doi:10.1016/j.euroneuro.2015.08.013
- Chen, Y., & Baram, T. Z. (2016). Toward understanding how early-life stress reprograms cognitive and emotional brain networks. *Neuropsychopharmacology*, *41*, 197-206. doi:10.1038/npp.2015.181
- Chou, C., Condrón, L., & Belland, J. C. (2005). A review of the research on Internet addiction. *Educational Psychology Review*, *17*, 363-387. doi:10.1007/s10648-005-8138-1
- Christiansen, P., Schoenmakers, T. M., & Field, M. (2015). Less than meets the eye: Reappraising the clinical relevance of attentional bias in addiction. *Addictive Behaviors*, *44*, 43–50.
- Ciccarelli, M., Nigro, G., Griffiths, M. D., Cosenza, M., & D'Olimpio, F. (2016). Attentional biases in problem and non-problem gamblers. *Journal of Affective Disorders*, *198*, 135–141.
- Correa, T., Hinsley, A. W., & de Zuniga, H. G. (2010). Who interacts on the Web? The intersection of users' personality and social media use. *Computers in Human Behavior*, *26*, 247-253.
- Cousijn, J., Goudriaan, A. E., Ridderinkhof, K. R., Van Den Brink, W., Veltman, D. J., & Wiers, R. W. (2012). Approach-bias predicts development of cannabis problem severity in heavy cannabis users: Results from a prospective fMRI study. *PLoS One*, *7*, e42394. doi:10.1371/journal.pone.0042394
- Cousijn, J., Goudriaan, A. E., & Wiers, R. W. (2011). Reaching out towards cannabis: Approach-bias in heavy cannabis users predicts changes in cannabis use. *Addiction*, *106*, 1667-1674. doi:10.1111/j.1360-0443.2011.03475.x
- Crockford, D. N., Goodyear, B., Edwards, J., Qickfall, J., & el-Guebaly, N. (2005). Cue-induced brain activity in pathological gamblers. *Biological Psychiatry*, *58*, 787-795.
- Czapla, M., Simon, J., Friederich, H.-C., Herpertz, S. C., Zimmermann, P., & Loeber, S. (2015). Is binge drinking in young adults associated with an alcohol-specific impairment of response inhibition? *European Addiction Research*, *21*, 105-113.

- Dalbudak, E., Evren, C., Aldemir, S., & Evren, B. (2014). The severity of Internet addiction risk and its relationship with the severity of borderline personality features, childhood traumas, dissociative experiences, depression and anxiety symptoms among Turkish university students. *Psychiatry Research*, *219*, 577–582.
- Davis, R. A. (2001). A cognitive-behavioral model of pathological Internet use. *Computers in Human Behavior*, *17*, 187-195. doi:10.1016/S0747-5632(00)00041-8
- Demetrovics, Z., Urbán, R., Nagygyörgy, K., Farkas, J., Zilahy, D., Mervó, B., . . . E., H. (2011). Why do you play? The development of the motives for online gaming questionnaire (MOGQ). *Behavior Research Methods*, *43*, 814–825. doi:10.3758/s13428-011-0091-y
- Derbyshire, K. L., & Grant, J. E. (2015). Compulsive sexual behavior: A review of the literature. *Journal of Behavioral Addictions*, *4*, 37-43. doi:10.1556/2006.4.2015.003
- Deryakulu, D., & Ursavas, Ö. F. (2014). Genetic and environmental influences on problematic Internet use: A twin study. *Computers in Human Behavior*, *39*, 331–338 doi:10.1016/j.chb.2014.07.038
- Dickerson, S. S., & Kemeny, M. E. (2004). Acute stressors and cortisol responses: A theoretical integration and synthesis of laboratory research. *Psychological Bulletin*, *130*, 355-391.
- Ding, W. N., Sun, J. H., Sun, Y. W., Chen, X., Zhou, Y., Zhuang, Z. G., . . . Du, Y. S. (2014). Trait impulsivity and impaired prefrontal impulse inhibition function in adolescents with internet gaming addiction revealed by a Go/No-Go fMRI study. *Behavioral and Brain Function*, *10*, 20. doi:10.1186/1744-9081-10-20
- Dong, G., Devito, E. E., Du, X., & Cui, Z. (2012). Impaired inhibitory control in “internet addiction disorder”: A functional magnetic resonance imaging study. *Psychiatry Research*, *203*, 153–158. doi:10.1016/j.psychres.2012.02.001
- Dong, G., Hu, Y., Lin, X., & Lu, Q. (2013). What makes Internet addicts continue playing online even when faced by severe negative consequences? Possible explanations from an fMRI study. *Biological Psychology*, *94*, 282-289. doi:10.1016/j.biopsycho.2013.07.009
- Dong, G., Lin, X., Hu, Y., Xie, C., & Du, X. (2015). Imbalanced functional link between executive control network and reward network explain the online-game seeking behaviors in Internet gaming disorder. *Scientific Reports*, *5*, 9197. doi:10.1038/srep09197
- Dong, G., Lin, X., Zhou, H., & Lu, Q. (2014). Cognitive flexibility in internet addicts: fMRI evidence from difficult-to-easy and easy-to-difficult switching situations. *Addictive Behaviors*, *39*, 677-683. doi:10.1016/j.addbeh.2013.11.028
- Dong, G., Lu, Q., Zhou, H., & Zhao, X. (2010). Impulse inhibition in people with Internet addiction disorder: Electrophysiological evidence from a Go/NoGo study. *Neuroscience Letters*, *485*, 138-142.
- Dong, G., & Potenza, M. N. (2014). A cognitive-behavioral model of Internet gaming disorder: Theoretical underpinnings and clinical implications. *Journal of Psychiatric Research*, *58*, 7-11. doi:10.1016/j.jpsychires.2014.07.005
- Dong, G., & Potenza, M. N. (2016). Risk-taking and risky decision-making in Internet gaming disorder: Implications regarding online gaming in the setting of negative consequences. *Journal of Psychiatric Research*, *73*, 1–8. doi:10.1016/j.jpsychires.2015.11.011

- Dong, G., Shen, Y., Huang, J., & Du, X. (2013). Impaired error-monitoring function in people with internet addiction disorder: An event-related fMRI study. *European Addiction Research, 19*, 269-275. doi:10.1159/000346783
- Dong, G., Zhou, H., & Zhao, X. (2011). Male Internet addicts show impaired executive control ability: Evidence from a color-word Stroop task. *Neuroscience Letters, 499*, 114–118. doi:0.1016/j.neulet.2011.05.047
- Douglas, A. C., Mills, J. E., Niang, M., Stepchenkova, S., Byun, S., Ruffini, C., . . . Blanton, M. (2008). Internet addiction: Meta-synthesis of qualitative research for the decade 1996-2006. *Computers in Human Behavior, 24*, 3027-3044.
- Duka, T., Trick, L., Nikolaou, K., Gray, M. A., Kempton, M. J., Williams, H., . . . Stephens, D. N. (2011). Unique brain areas associated with abstinence control are damaged in multiply detoxified alcoholics. *Biological Psychiatry, 70*, 545-552. doi:10.1016/j.biopsych.2011.04.006
- Ebeling-Witte, S., Frank, M. L., & Lester, D. (2007). Shyness, Internet use, and personality. *Cyberpsychology & Behavior, 10*, 713-716. doi:10.1089/cpb.2007.9964
- Eberl, C., Wiers, R. W., Pawelczack, S., Rinck, M., Becker, E. S., & Lindenmeyer, J. (2013a). Approach bias modification in alcohol dependence: Do clinical effects replicate and for whom does it work best? *Developmental Cognitive Neuroscience, 4*, 38-51. doi:10.1016/j.dcn.2012.11.002
- Eberl, C., Wiers, R. W., Pawelczack, S., Rinck, M., Becker, E. S., & Lindenmeyer, J. (2013b). Implementation of approach bias re-training in alcoholism. How many sessions are needed? *Alcoholism: Clinical and Experimental Research, 38*(2), 587-594. doi:10.1111/acer.12281
- Elsley, J., Coates, A., Lacadie, C. M., McCrory, E. J., Sinha, R., Mayes, L. C., & Potenza, M. N. (2015). Childhood trauma and neural responses to personalized stress, favorite-food and neutral-relaxing cues in adolescents. *Neuropsychopharmacology, 40*, 1580-1589.
- Evans, J. S. B. T. (2003). In two minds: Dual-process accounts of reasoning. *Trends in Cognitive Sciences, 7*, 454-459. doi:10.1016/j.tics.2003.08.012
- Evans, J. S. B. T., & Coventry, K. (2006). A dual process approach to behavioural addiction: The case of gambling. In R. W. Wiers & A. W. Stacy (Eds.), *Handbook of Implicit Cognition and Addiction* (pp. 29–43). Thousand Oaks, CA: Sage.
- Everitt, B. J. (2014). Neural and psychological mechanisms underlying compulsive drug seeking habits and drug memories - indications for novel treatments of addiction. *European Journal of Neuroscience, 40*, 2163-2182.
- Everitt, B. J., & Robbins, T. W. (2005). Neural systems of reinforcement for drug addiction: From actions to habits to compulsion. *Nature Neuroscience, 8*, 1481–1489. doi:10.1038/nn1579
- Everitt, B. J., & Robbins, T. W. (2016). Drug addiction: Updating actions to habits to compulsions ten years on. *Annual Review of Psychology, 67*, 23-50. doi:10.1146/annurev-psych-122414-033457
- Fauth-Bühler, M., & Mann, K. (2015). Neurobiological correlates of internet gaming disorder: Similarities to pathological gambling. *Addictive Behaviors, EPub ahead of print*. doi:10.1016/j.addbeh.2015.11.004
- Fauth-Bühler, M., Mann, K., & Potenza, M. N. (2016). Pathological gambling: A review of the neurobiological evidence relevant for its classification as an addictive disorder. *Addiction Biology*. doi:10.1111/adb.12378

- Field, M., & Cox, W. M. (2008). Attentional bias in addictive behaviors: A review of its development, causes, and consequences. *Drug and Alcohol Dependence*, *97*, 1-20. doi:10.1016/j.drugalcdep.2008.03.030
- Field, M., Munafò, M. R., & Franken, I. H. A. (2009). A meta-analytic investigation of the relationship between attentional bias and subjective craving in substance abuse. *Psychological Bulletin*, *135*, 589-607. doi:10.1037/a0015843
- Floros, G., Siomos, K., Stogiannidou, A., Giouzepas, I., & Garyfallos, G. (2014). The relationship between personality, defense styles, internet addiction disorder, and psychopathology in college students. *Cyberpsychology, Behavior and Social Networking*, *17*, 672-676.
- Forrest, L. N., Smith, A. R., Fussner, L. M., Dodd, D. R., & Clerkin, E. M. (2016). Using implicit attitudes of exercise importance to predict explicit exercise dependence symptoms and exercise behaviors. *Psychology of Sport and Exercise*, *22*, 91-97. doi:10.1016/j.psychsport.2015.06.006
- Goldstein, R. Z., Craig, A. D., Bechara, A., Garavan, H., Childress, A. R., Paulus, M. P., & Volkow, N. D. (2009). The neurocircuitry of impaired insight in drug addiction. *Trends in Cognitive Sciences*, *13*, 372-380. doi:10.1016/j.tics.2009.06.004
- Goldstein, R. Z., & Volkow, N. D. (2002). Drug addiction and its underlying neurobiological basis: Neuroimaging evidence for the involvement of the frontal cortex. *American Journal of Psychiatry*, *159*, 1642-1652. doi:10.1176/appi.ajp.159.10.1642
- Goldstein, R. Z., & Volkow, N. D. (2011). Dysfunction of the prefrontal cortex in addiction: Neuroimaging findings and clinical implications. *Nature Reviews Neuroscience*, *12*, 652-669.
- Goudriaan, A. E., de Ruiter, M. B., van den Brink, W., Oosterlaan, J., & Veltman, D. J. (2010). Brain activation patterns associated with cue reactivity and craving in abstinent problem gamblers, heavy smokers and healthy controls: An fMRI study. *Addiction Biology*, *15*, 491-503. doi:10.1111/j.1369-1600.2010.00242.x
- Goudriaan, A. E., Oosterlaan, J., Beurs, E., & van den Brink, W. (2004). Pathological gambling: A comprehensive review of biobehavioral findings. *Neuroscience and Biobehavioral Reviews*, *28*, 123-141. doi:10.1016/j.neubiorev.2004.03.001
- Grant, J. E., Brewer, J. A., & Potenza, M. N. (2006). The neurobiology of substance and behavioral addictions. *CNS Spectrum*, *11*, 924-930.
- Grant, J. E., & Chamberlain, S. R. (2015). Gambling disorder and its relationship with substance use disorders: Implications for nosological revisions and treatment. *American Journal of Addiction*, *24*, 126-131. doi:10.1111/ajad.12112
- Grant, J. E., Potenza, M. N., Weinstein, A., & Gorelick, D. A. (2010). Introduction to behavioral addictions. *The American Journal of Drug and Alcohol Abuse*, *36*, 233-241.
- Grant, J. E., Schreiber, L. R., & Odlaug, B. L. (2013). Phenomenology and treatment of behavioural addictions. *Canadian Journal of Psychiatry*, *58*, 252-259.
- Greenwald, A. G., McGhee, D. E., & Schwartz, J. L. (1998). Measuring individual differences in implicit cognition: The implicit association test. *Journal of Personality and Social Psychology*, *74*, 1464-1480.
- Griffiths, M. D. (1999). Internet addiction: Fact or fiction? *Psychologist*, *12*, 246-250.
- Griffiths, M. D. (2012). Internet sex addiction: A review of empirical research. *Addiction Research and Theory*, *20*, 111-124. doi:10.3109/16066359.2011.588351

- Gross, J. J., & Jazaieri, H. (2014). Emotion, emotion regulation, and psychopathology: An affective science perspective. *Clinical Psychological Science*, 2, 387–401. doi:10.1177/2167702614536164
- Han, D. H., Lee, Y. S., Yang, K. C., Kim, E. Y., Lyoo, I. K., & Renshaw, P. F. (2007). Dopamine genes and reward dependence in adolescents with excessive Internet video game play. *Journal of Addiction Medicine*, 1, 133-138.
- Hansen, S. (2002). Excessive Internet usage or 'Internet Addiction'? The implications of diagnostic categories for student users. *Journal of Computer Assisted Learning*, 18, 235-236. doi:10.1046/j.1365-2729.2002.t01-2-00230.x
- Hardie, E., & Tee, M. Y. (2007). Excessive Internet use: The role of personality, loneliness, and social support networks in Internet Addiction. *Australian Journal of Emerging Technologies and Society*, 5, 34-47.
- Ho, R. C., Zhang, M. W. B., Tsang, T. Y., Toh, A. H., Pan, F., Lu, Y., . . . Mak, K.-K. (2014). The association between internet addiction and psychiatric co-morbidity: A meta-analysis. *BMC Psychiatry*, 14, 183. doi:10.1186/1471-244X-14-183
- Hogarth, L., Dickinson, A., & Duka, T. (2010). The associative basis of cue-elicited drug taking in humans. *Psychopharmacology*, 208, 337-351. doi:10.1007/s00213-009-1735-9
- Hogarth, L., Dickinson, A., Hutton, S. B., Elbers, N., & Duka, T. (2006). Drug expectancy is necessary for stimulus control of human attention, instrumental drug-seeking behaviour and subjective pleasure. *Psychopharmacology*, 185, 495-504. doi:10.1007/s00213-005-0287-x
- Hogarth, L., Dickinson, A., Wright, A., Kouvaraki, M., & Duka, T. (2007). The role of drug expectancy in the control of human drug seeking. *Journal of Experimental Psychology: Animal Behavior Processes*, 33, 484-496. doi:10.1037/0097-7403.33.4.484
- Holahan, C. J., Moos, R. H., Holahan, C. K., Cronkite, R. C., & Randall, P. K. (2001). Drinking to cope, emotional distress and alcohol use and abuse: A ten-year model. *Journal of Studies on Alcohol*, 62, 190–198. doi:10.15288/jsa.2001.62.190
- Hong, S.-B., Kim, J.-W., Choi, E.-J., Kim, H.-H., Suh, J.-E., Kim, C.-D., . . . Yi, S.-H. (2013). Reduced orbitofrontal cortical thickness in male adolescents with internet addiction. *Behavioral and Brain Function*, 9, 11. doi:10.1186/1744-9081-9-11
- Hong, S.-B., Zalesky, A., Cocchi, L., Fornito, A., Choi, E.-J., Kim, H.-H., . . . Yi, S.-H. (2013). Decreased functional brain connectivity in adolescents with internet addiction. *PLoS One*, 8, e57831. doi:10.1371/journal.pone.0057831
- Hormes, J. M., Kearns, B., & Timko, C. A. (2014). Craving Facebook? Behavioral addiction to online social networking and its association with emotion regulation deficits. *Addiction*, 109, 2079–2088. doi:10.1111/add.12713
- Houben, K., & Jansen, A. (2011). Training inhibitory control. A recipe for resisting sweet temptations. *Appetite*, 56, 345-349. doi:10.1016/j.appet.2010.12.017
- Houben, K., Nederkoorn, C., Wiers, R. W., & Jansen, A. (2011). Resisting temptation: Decreasing alcohol-related affect and drinking behavior by training response inhibition. *Drug and Alcohol Dependence*, 116, 132-136. doi:10.1016/j.drugalcdep.2010.12.011
- Houben, K., Schoenmakers, T. M., & Wiers, R. W. (2010). I didn't feel like drinking but I don't know why: The effects of evaluative conditioning on alcohol-related attitudes, craving and behavior. *Addictive Behaviors*, 35(12), 1161-1163. doi:10.1016/j.addbeh.2010.08.012

- Hsieh, Y. P., Shen, A. C. T., Wei, H. S., Feng, J. Y., Huang, S. C. Y., & Hwa, H. L. (2016). Associations between child maltreatment, PTSD, and internet addiction among Taiwanese students. *Computers in Human Behavior*, *56*, 209–214. doi:10.1016/j.chb.2015.11.048
- Jeromin, F., Nyenhuis, N., & Barke, A. (2016). Attentional bias in excessive Internet gamers: Experimental investigations using an addiction Stroop and a visual probe. *Journal of Behavioral Addictions*, *5*, 32-40.
- Jović, J., & Đinđić, N. (2011). Influence of dopaminergic system on Internet addiction. *Acta Medica Medianae*, *50*, 60-66. doi:10.5633/amm.2011.0112
- Kahneman, D. (2003). A perspective on judgment and choice: Mapping bounded rationality. *American Psychologist*, *58*, 697-720. doi:10.1037/0003-066X.58.9.697
- Kalivas, P. W., & Volkow, N. D. (2005). The neural basis of addiction: A pathology of motivation and choice. *American Journal of Psychiatry*, *162*, 1403–1413.
- Kaptsis, D., King, D. L., Delfabbro, P. H., & Gradisar, M. (2016). Withdrawal symptoms in internet gaming disorder: A systematic review. *Clinical Psychology Review*, *43*, 58-66. doi:10.1016/j.cpr.2015.11.006
- Kardefelt-Winther, D. (2014). A conceptual and methodological critique of internet addiction research: Towards a model of compensatory internet use. *Computers in Human Behavior*, *31*, 351–354. doi:10.1016/j.chb.2013.10.059
- Kim, H. K., & Davis, K. E. (2009). Toward a comprehensive theory of problematic Internet use: Evaluating the role of self-esteem, anxiety, flow, and the self-rated importance of Internet activities. *Computers in Human Behavior*, *25*, 490-500. doi:10.1016/j.chb.2008.11.001
- Kim, S. H., Baik, S.-H., Park, C. S., Kim, S. J., Choi, S. W., & Kim, S. E. (2011). Reduced striatal dopamine D2 receptors in people with Internet addiction. *Neuroreport*, *22*, 407-411. doi:10.1097/WNR.0b013e328346e16e
- King, D. L., & Delfabbro, P. H. (2014). The cognitive psychology of Internet gaming disorder. *Clinical Psychology Review*, *34*, 298–308.
- Klucken, T., Wehrum-Osinsky, S., Schweckendiek, J., Kruse, O., & Stark, R. (2016). Altered appetitive conditioning and neural connectivity in subjects with compulsive sexual behavior. *Journal of Sexual Medicine*, *13*, 627-636. doi:10.1016/j.jsxm.2016.01.013
- Ko, C.-H., Liu, G. C., Hsiao, S., Yen, J. Y., Yang, M. J., Lin, W. C., . . . Chen, C. S. (2009). Brain activities associated with gaming urge of online gaming addiction. *Journal of Psychiatric Research*, *43*, 739-747. doi:10.1016/j.jpsychires.2008.09.012
- Kober, H., Lacadie, C., Wexler, B. E., Malison, R. T., Sinha, R., & Potenza, M. N. (2016). Brain activity during cocaine craving and gambling urges: An fMRI study. *Neuropsychopharmacology*, *41*, 628-637.
- Koo, H. J., & Kwon, J. H. (2014). Risk and protective factors of Internet addiction: A meta-analysis of empirical studies in Korea. *Yonsei Medical Journal*, *55*, 1691-1711.
- Koob, G. F., & Volkow, N. D. (2010). Neurocircuitry of addiction. *Neuropsychopharmacology*, *35*, 217–238.
- Koolhaas, J. M., Bartolomucci, A., Buwalda, B., de Boer, S. F., Flügge, G., Korte, S. M., . . . Fuchs, E. (2011). Stress revisited: A critical evaluation of the stress concept. *Neuroscience & Biobehavioral Reviews*, *35*, 1291-1301. doi:10.1016/j.neubiorev.2011.02.003

- Kor, A., Zilcha-Mano, S., Fogel, Y. A., Mikulincer, M., Reid, R. C., & Potenza, M. N. (2014). Psychometric development of the Problematic Pornography Use Scale. *Addictive Behaviors*, *39*, 861–868. doi:10.1016/j.addbeh.2014.01.027
- Kraus, S. W., Voon, V., & Potenza, M. N. (2016). Should compulsive sexual behavior be considered an addiction? *Addiction*, *Epub ahead of print*. doi:10.1111/add.13297
- Kukar-Kinney, M., Ridgway, N. M., & Monroe, K. B. (2009). The relationship between consumers' tendencies to buy compulsively and their motivations to shop and buy on the Internet. *Journal of Retailing*, *85*, 298-307.
- Kuss, D. J. (2013). Internet gaming addiction: Current perspectives. *Psychology Research and Behavior Management*, *6*, 125–137.
- Kuss, D. J., & Griffiths, M. D. (2011a). Internet gaming addiction: A systematic review of empirical research. *International Journal Of Mental Health And Addiction*, *10*, 278-296. doi:0.1007/s11469-011-9318-5
- Kuss, D. J., & Griffiths, M. D. (2011b). Online social networking and addiction: A review of the psychological literature. *International Journal of Enviromental Research and Public Health*, *8*, 3528-3552. doi:10.3390/ijerph8093528
- Kuss, D. J., & Griffiths, M. D. (2012). Internet and gaming addiction: A systematic literature review of neuroimaging studies. *Brain Sciences*, *2*, 347-374. doi:10.3390/brainsci2030347
- Kuss, D. J., Griffiths, M. D., Karila, M., & Billieux, J. (2014). Internet addiction: A systematic review of epidemiological research for the last decade. *Current Pharmaceutical Design*, *20*, 4026-4052.
- Kuss, D. J., & Lopez-Fernandez, O. (2016). Internet addiction and problematic Internet use: A systematic review of clinical research. *World Journal of Psychiatry*, *6*, 143–176. doi:10.5498/wjp.v6.i1.143.
- Kuss, D. J., Louws, J., & Wiers, R. W. (2012). Online gaming addiction? Motives predict addictive play behavior in Massively Multiplayer Online Role-Playing Games. *Cyberpsychology, Behavior and Social Networking*, *15*, 480–485.
- Laier, C., & Brand, M. (2014). Empirical evidence and theoretical considerations on factors contributing to cybersex addiction from a cognitive-behavioral view. *Sexual Addiction & Compulsivity*, *21*, 305-321. doi:10.1080/10720162.2014.970722
- Laier, C., Pawlikowski, M., & Brand, M. (2014). Sexual picture processing interferes with decision-making under ambiguity. *Archives of Sexual Behavior*, *43*, 473-482. doi:10.1007/s10508-013-0119-8
- Laier, C., Pawlikowski, M., Pekal, J., Schulte, F. P., & Brand, M. (2013). Cybersex addiction: Experienced sexual arousal when watching pornography and not real-life sexual contacts makes the difference. *Journal of Behavioral Addictions*, *2*, 100-107. doi:10.1556/JBA.2.2013.002
- Lam, L. T., & Wong, E. M. (2015). Stress moderates the relationship between problematic Internet use by parents and problematic Internet use by adolescents. *Journal of Adolescent Health*, *56*, 300-306. doi:10.1016/j.jadohealth.2014.10.263
- Lee, S., & Lee, J.-H. (2015). The effect of automatic attentional bias modification on alcohol ambivalence. *Addictive Behaviors*, *46*, 58-64. doi:10.1016/j.addbeh.2015.03.010
- Lee, Y. H., Ko, C. H., & Chou, C. (2014). Re-visiting Internet addiction among Taiwanese students: A cross-sectional comparison of students' expectations, online gaming, and online social interaction. *Journal of Abnormal Child Psychology*, [Epub ahead of print].

- Lee, Y. S., Han, D. H., Yang, K. C., Daniels, M. A., Na, C., Kee, B. S., & Renshaw, P. F. (2008). Depression like characteristics of 5HTTLPR polymorphism and temperament in excessive internet users. *Journal of Affective Disorders, 109*, 165–169.
- Li, J.-R., Wang, C.-H., & Lin, C.-W. (2012). Locus of control, emotion venting strategy and Internet addiction. *International Journal of Social, Behavioral, Educational, Economic, Business and Industrial Engineering, 6*, 3392-3395.
- Li, M., Chen, J., Li, N., & Li, X. (2014). A twin study of problematic internet use: Its heritability and genetic association with effortful control. *Twin Research and Human Genetics, 17*, 279–287.
- Liu, L., Yip, S. W., Zhang, J.-T., Wang, L.-J., Shen, Z.-J., Liu, B., . . . Fang, X. Y. (2016). Activation of the ventral and dorsal striatum during cue reactivity in Internet gaming disorder. *Addiction Biology*.
- Loeber, S., & Duka, T. (2009). Acute alcohol impairs conditioning of a behavioural reward-seeking response and inhibitory control processes - Implications for addictive disorders. *Addiction, 104*, 2013–2022. doi:10.1111/j.1360-0443.2009.02718.x
- Lu, H., Ma, L., Lee, T., Hou, H., & Liao, H. (2014). The link of sexual sensation seeking to acceptance of cybersex, multiple sexual partners, and one-night stands among Taiwanese college students. *Journal of Nursing Research, 22*, 208-215.
- Luijten, M., Meerkerk, G. J., Franken, I. H., van de Wetering, B. J., & Schoenmakers, T. M. (2015). An fMRI study of cognitive control in problem gamers. *Psychiatry Research, 231*, 262-268. doi:10.1016/j.psychresns.2015.01.004
- Mechelmans, D. J., Irvine, M., Banca, P., Porter, L., Mitchell, S., Mole, T. B., . . . Voon, V. (2014). Enhanced attentional bias towards sexually explicit cues in individuals with and without compulsive sexual behaviours. *PLoS One, 9*, e105476.
- Meng, Y., Deng, W., Wang, H., Guo, W., & Li, T. (2015). The prefrontal dysfunction in individuals with Internet gaming disorder: A meta-analysis of functional magnetic resonance imaging studies. *Addiction Biology, 20*, 799–808. doi:10.1111/adb.12154
- Metcalfe, O., & Pammer, K. (2011). Attentional bias in excessive massively multiplayer online role-playing gamers using a modified stroop task. *Computers in Human Behavior, 27*, 1942-1947. doi:10.1016/j.chb.2011.05.001
- Miedl, S. F., Büchel, C., & Peters, J. (2014). Cue-induced craving increases impulsivity via changes in striatal value signals in problem gamblers. *Journal of Neuroscience, 34*, 4750-4755. doi:10.1523/JNEUROSCI.5020-13.2014
- Montag, C., Bey, K., Sha, P., Li, M., Chen, Y. F., Liu, W. Y., . . . Reuter, M. (2015). Is it meaningful to distinguish between generalized and specific Internet addiction? Evidence from a cross-cultural study from Germany, Sweden, Taiwan and China. *Asia-Pacific Psychiatry, 7*, 20-26.
- Montag, C., Kirsch, P., Sauer, C., Markett, S., & Reuter, M. (2012). The role of the CHRNA4 gene in internet addiction: A case-control Study. *Journal of Addiction Medicine, 6*, 191–195
- Morahan-Martin, J., & Schumacher, P. (2003). Loneliness and social uses of the internet. *Computers in Human Behavior, 19*, 659-671. doi:10.1016/S0747-5632(03)00040-2
- Müller, A., Brand, M., Mitchell, J. E., & de Zwaan, M. (in press). Pathological Online Shopping. In M. Potenza (Ed.), *Online Addiction*. Oxford: Oxford University Press.

- Müller, K. W., Beutel, M. E., Egloff, B., & Wöfling, K. (2014). Investigating risk factors for Internet Gaming Disorder: A comparison of patients with addictive gaming, pathological gamblers and healthy controls regarding the Big Five personality traits. *European Addiction Research, 20*, 129-136. doi:10.1159/000355832
- Müller, K. W., Dreier, M., Beutel, M., Duven, E., Giral, S., & Wöfling, K. (2016). A hidden type of internet addiction? Intense and addictive use of social networking sites in adolescents. *Computers in Human Behavior, 55*, 172-177. doi:10.1016/j.chb.2015.09.007
- Nie, J., Zhang, W., Chen, J., & Li, W. (2016). Impaired inhibition and working memory in response to internet-related words among adolescents with internet addiction: A comparison with attention-deficit/hyperactivity disorder. *Psychiatry Research, 236*, 28-34. doi:10.1016/j.psychres.2016.01.004
- Niemz, K., Griffiths, M. D., & Banyard, P. (2005). Prevalence of pathological Internet use among university students and correlations with self-esteem, the General Health Questionnaire (GHQ), and disinhibition. *Cyberpsychology & Behavior, 8*, 562-570. doi:10.1089/cpb.2005.8.562
- Noh, D., & Kim, S. (2016). Dysfunctional attitude mediates the relationship between psychopathology and Internet addiction among Korean college students: A cross-sectional observational study. *International Journal of Mental Health Nursing, 25*(12), 1222-1230. doi:10.1111/inm.12220
- Odacı, H., & Çikrikçi, Ö. (2014). Problematic internet use in terms of gender, attachment styles and subjective well-being in university students. *Computers in Human Behavior, 32*, 61-66.
- Odacı, H., & Kalkan, M. (2010). Problematic Internet use, loneliness and dating anxiety among young adult university students. *Computers and Education, 55*, 1091-1097.
- Osborne, L. A., Romano, M., Re, F., Roaro, A., Truzoli, R., & Reed, P. (2016). Evidence for an Internet addiction disorder: Internet exposure reinforces color preference in withdrawn problem users. *Journal of Clinical Psychiatry, 77*, 269-274. doi:10.4088/JCP.14m09658
- Park, C.-B., Park, S. M., Gwak, A. R., Sohn, B. K., Lee, J.-Y., Jung, H. Y., . . . Choi, J.-S. (2015). The effect of repeated exposure to virtual gambling cues on the urge to gamble. *Addictive Behaviors, 41*, 61-64. doi:10.1016/j.addbeh.2014.09.027
- Paul, B., & Shim, J. W. (2008). Gender, sexual affect, and motivations for Internet pornography use. *International Journal of Sexual Health, 20*, 187-199. doi:10.1080/19317610802240154
- Pawlikowski, M., & Brand, M. (2011). Excessive Internet gaming and decision making: Do excessive World of Warcraft-players have problems in decision making under risky conditions? *Psychiatry Research, 188*, 428-433. doi:10.1016/j.psychres.2011.05.017
- Pawlikowski, M., Nader, I. W., Burger, C., Biermann, I., Stieger, S., & Brand, M. (2014). Pathological Internet use – It is a multidimensional and not a unidimensional construct. *Addiction Research & Theory, 22*, 166-175. doi:10.3109/16066359.2013.793313
- Pericot-Valverde, I., García-Rodríguez, O., Gutiérrez-Maldonado, J., & Secades-Villa, R. (2015). Individual variables related to craving reduction in cue exposure treatment. *Addictive Behaviors, 49*, 59-63. doi:10.1016/j.addbeh.2015.05.013
- Petry, N. M., & O'Brian, C. P. (2013). Internet gaming disorder and the DSM-5. *Addiction, 108*, 1186-1187. doi:10.1111/add.12162

- Pezoa-Jares, R. E., Espinoza-Luna, I. L., & Vasquez-Medina, J. A. (2012). Internet addiction: A review. *Journal of Addiction Research & Therapy*, *S6*, 004. doi:10.4172/2155-6105.S6-004
- Piazza, P. V., & Deroche-Gamonet, V. (2013). A multistep general theory of transition to addiction. *Psychopharmacology*, *229*, 387-413.
- Pike, E., Stoops, W. W., Fillmore, M. T., & Rush, C. R. (2013). Drug-related stimuli impair inhibitory control in cocaine abusers. *Drug and Alcohol Dependence*, *133*, 768-771. doi:10.1016/j.drugalcdep.2013.08.004
- Pontes, H. M., Griffiths, M. D., & Patrão, I. M. (2014). Internet addiction and loneliness among children and adolescents in the education setting: An empirical pilot study. *Aloma*, *32*(91-98).
- Pontes, H. M., Kuss, D. J., & Griffiths, M. D. (2015). Clinical psychology of Internet addiction: A review of its conceptualization, prevalence, neuronal processes, and implications for treatment. *Neuroscience and Neuroeconomics*, *4*, 11–23.
- Potenza, M. N. (2006). Should addictive disorders include non-substance-related conditions? *Addiction*, *101*, 142-151. doi:10.1111/j.1360-0443.2006.01591.x
- Potenza, M. N. (2008). The neurobiology of pathological gambling and drug addiction: An overview and new findings. *Philosophical Transactions of the Royal Society of London Series B*, *363*, 3181–3189.
- Potenza, M. N. (2013). How central is dopamine to pathological gambling or gambling disorder? *Frontiers in Behavioral Neuroscience*, *7*, 206. doi:10.3389/fnbeh.2013.00206
- Potenza, M. N., Steinberg, M. A., Skudlarski, P., Fulbright, R. K., Lacadie, C. M., Wilber, M. K., . . . Wexler, B. E. (2003). Gambling urges in pathological gambling. A functional magnetic resonance imaging study. *Archives of General Psychiatry*, *60*, 828-836.
- Prizant-Passal, S., Shechner, T., & Aderka, I. M. (2016). Social anxiety and internet use – A meta-analysis: What do we know? What are we missing? *Computers in Human Behavior*, *62*, 221–229. doi:10.1016/j.chb.2016.04.003
- Reid, R. C., Garos, S., & Carpenter, B. N. (2011). Reliability, validity, and psychometric development of the Hypersexual Behavior Inventory in an outpatient sample of men. *Sexual Addiction & Compulsivity*, *18*, 30–51.
- Rinck, M., & Becker, E. (2007). Approach and avoidance in fear of spiders. *Journal of Behavior Therapy and Experimental Psychiatry*, *38*, 105-120.
- Robbins, T. W., & Clark, L. (2015). Behavioral addictions. *Current Opinions in Neurobiology*, *30*, 66-72. doi:10.1016/j.conb.2014.09.005
- Robinson, T. E., & Berridge, K. C. (1993). The neural basis of drug craving: An incentive-sensitization theory of addiction. *Brain research. Brain research reviews*, *18*, 247-291.
- Robinson, T. E., & Berridge, K. C. (2000). The psychology and neurobiology of addiction: An incentive–sensitization view. *Addiction*, *95*, 91-117.
- Robinson, T. E., & Berridge, K. C. (2001). Incentive-sensitization and addiction. *Addiction*, *96*, 103-114.
- Robinson, T. E., & Berridge, K. C. (2008). The incentive sensitization theory of addiction: Some current issues. *Philosophical Transactions of the Royal Society B*, *363*, 3137-3146. doi:10.1098/rstb.2008.0093
- Romano, M., Osborne, L. A., Truzoli, R., & Reed, P. (2013). Differential psychological impact of internet exposure on internet addicts. *PLoS One*, *8*, e55162. doi:10.1371/journal.pone.0055162

- Rooke, S. E., Hine, D. W., & Thorsteinsson, E. B. (2008). Implicit cognition and substance use: A meta-analysis. *Addictive Behaviors, 33*, 1314-1328. doi:10.1016/j.addbeh.2008.06.009
- Ryan, R. M., Rigby, C. S., & Przybylski, A. (2006). The motivational pull of video games: A self-determination theory approach. *Motivation and Emotion, 30*, 347-363.
- Ryan, T., & Xenos, S. (2011). Who uses Facebook? An investigation into the relationship between the Big Five, shyness, narcissism, loneliness, and Facebook usage. *Computers in Human Behavior, 27*, 1658-1664.
- Santos, V. A., Freire, R., Zugliani, M., Cirillo, P., Santos, H. H., Nardi, A. E., & King, A. L. (2016). Treatment of Internet addiction with anxiety disorders: Treatment protocol and preliminary before-after results involving pharmacotherapy and modified cognitive behavioral therapy. *JMIR Research Protocols, 5*, e46. doi:10.2196/resprot.5278
- Sariyska, R., Reuter, M., Bey, K., Sha, P., Li, M., Chen, Y. F., . . . Montag, C. (2014). Self-esteem, personality and Internet addiction: A cross-cultural comparison study. *Personality and Individual Differences, 61-62*, 28-33.
- Sarnyai, Z., & Kovács, G. L. (2014). Oxytocin in learning and addiction: From early discoveries to the present. *Pharmacology Biochemistry and Behavior, 119*, 3–9. doi:10.1016/j.pbb.2013.11.019
- Schiebener, J., & Brand, M. (2015). Decision making under objective risk conditions - a review of cognitive and emotional correlates, strategies, feedback processing, and external influences. *Neuropsychology Review, 25*, 171-198.
- Schimmenti, A., Passanisi, A., Gervasi, A. M., Manzella, S., & Famà, F. I. (2014). Insecure attachment attitudes in the onset of problematic internet use among late adolescents. *Child Psychiatry and Human Development, 45*, 588–595.
- Schoenmakers, T. M., de Bruin, M., Lux, I. F., Goertz, A. G., van Kerkhof, D. H., & Wiers, R. W. (2010). Clinical effectiveness of attentional bias modification training in abstinent alcoholic patients. *Drug and Alcohol Dependence, 109*, 30-36. doi:10.1016/j.drugalcdep.2009.11.022
- Seok, J. W., Lee, K. H., Sohn, S., & Sohn, J. H. (2015). Neural substrates of risky decision making in individuals with Internet addiction. *Australian and New Zealand Journal of Psychiatry, 49*, 923-932. doi:10.1177/0004867415598009
- Sepede, G., Tavino, M., Santacroce, R., Fiori, F., Salerno, R. M., & Di Giannantonio, M. (2016). Functional magnetic resonance imaging of internet addiction in young adults. *World Journal of Radiology, 8*, 210–225. doi:10.4329/wjr.v8.i2.210
- Shapiro, D., Jamner, L. D., Davydov, D. M., & James, P. (2002). Situations and moods associated with smoking in everyday life. *Psychology of Addictive Behaviors, 16*, 342–345. doi:10.1037/0893-164X.16.4.342
- Smith, J. L., Mattick, R. P., Jamadar, S. D., & Iredale, J. M. (2014). Deficits in behavioural inhibition in substance abuse and addiction: a meta-analysis. *Drug and Alcohol Dependence, 145*, 1-33. doi:10.1016/j.drugalcdep.2014.08.009
- Snagowski, J., & Brand, M. (2015). Symptoms of cybersex addiction can be linked to both approaching and avoiding pornographic stimuli: Results from an analog sample of regular cybersex users. *Frontiers in Psychology, 6*, 653.
- Snagowski, J., Laier, C., Duka, T., & Brand, M. (in press). Subjective craving for pornography and associative learning predict tendencies towards cybersex addiction in a sample of regular cybersex users. *Sexual Addiction & Compulsivity*.

- Snagowski, J., Wegmann, E., Pekal, J., Laier, C., & Brand, M. (2015). Implicit associations in cybersex addiction: Adaption of an Implicit Association Test with pornographic pictures. *Addictive Behaviors*, *49*, 7–12.
- Spada, M. M. (2014). An overview of problematic Internet use. *Addictive Behaviors*, *39*, 3-6. doi:10.1016/j.addbeh.2013.09.007
- Stacy, A. W., & Wiers, R. W. (2010). Implicit cognition and addiction: A tool for explaining paradoxical behavior. *Annual Review of Clinical Psychology*, *6*, 551-575. doi:10.1146/annurev.clinpsy.121208.131444
- Stanovich, K. E., & West, R. F. (2000). Individual differences in reasoning: Implications for the rationality debate? *Behavioral and Brain Sciences*, *23*, 645-726.
- Starcevic, V. (2013). Is Internet addiction a useful concept? *Australian and New Zealand Journal of Psychiatry*, *47*, 16-19. doi:10.1177/0004867412461693
- Starcke, K., & Brand, M. (2012). Stress and decision making: A selective review. *Neuroscience & Biobehavioral Reviews*, *36*, 1228-1248. doi:10.1016/j.neubiorev.2012.02.003
- Starcke, K., & Brand, M. (in press). Effects of stress on decisions under uncertainty: A meta-analysis. *Psychological Bulletin*.
- Strittmatter, E., Parzer, P., Brunner, R., Fischer, G., Durkee, T., Carli, V., . . . Kaess, M. (2016). A 2-year longitudinal study of prospective predictors of pathological Internet use in adolescents. *European Child & Adolescence Psychiatry*, *25*, 725-734. doi:10.1007/s00787-015-0779-0
- Suissa, A. J. (2015). Cyber addictions: Toward a psychosocial perspective. *Addictive Behaviors*, *43*, 28-32. doi:10.1016/j.addbeh.2014.09.020
- Sun, D.-L., Chen, Z. J., Ma, N., Zhang, X.-C., Fu, X.-M., & Zhang, D. R. (2009). Decision-making and prepotent response inhibition functions in excessive Internet users. *CNS Spectrums*, *14*, 75-81.
- Tang, J., Yu, Y., Du, Y., Ma, Y., Zhang, D., & Wang, J. (2014). Prevalence of internet addiction and its association with stressful life events and psychological symptoms among adolescent internet users. *Addictive Behaviors*, *39*, 744-747. doi:10.1016/j.addbeh.2013.12.010
- Tavolacci, M., Ladner, J., Grigioni, S., Richard, L., Villet, H., & Dechelotte, P. (2013). Prevalence and association of perceived stress, substance use and behavioral addictions: A cross-sectional study among university students in France, 2009–2011. *BMC Public Health*, *13*, 724.
- Taymur, I., Budak, E., Demirci, H., Akdağ, H. A., Güngör, B. B., & Özdel, K. (2016). A study of the relationship between internet addiction, psychopathology and dysfunctional beliefs. *Computers in Human Behavior*, *61*, 532–536.
- Thalemann, R., Wöfling, K., & Grüsser, S. M. (2007). Specific cue reactivity on computer game-related cues in excessive gamers. *Behavioral Neuroscience*, *121*, 614-618. doi:10.1037/0735-7044.121.3.614
- Thatcher, A., Wretschko, G., & Fridjhon, P. (2008). Online flow experiences, problematic Internet use and Internet procrastination. *Computers in Human Behavior*, *24*, 2236-2254. doi:10.1016/j.chb.2007.10.008
- Thorberg, F. A., & Lyvers, M. (2006). Negative mood regulation (NMR) expectancies, mood, and affect intensity among clients in substance disorder treatment facilities. *Addictive Behaviors*, *31*, 811–820. doi:10.1016/j.addbeh.2005.06.008
- Tiffany, S. T., Carter, B. L., & Singleton, E. G. (2000). Challenges in the manipulation, assessment and interpretation of craving relevant variables. *Addiction*, *95*, 177-187.

- Tonioni, F., Mazza, M., Autullo, G., Cappelluti, R., Catalano, V., Marano, G., . . . Lai, C. (2014). Is Internet addiction a psychopathological condition distinct from pathological gambling? *Addictive Behaviors*, *39*, 1052-1056. doi:10.1016/j.addbeh.2014.02.016
- Turel, O., Serenko, A., & Giles, P. (2011). Integrating technology addiction and use: An empirical investigation of online auction users. *MIS Quarterly*, *35*, 1043-1061.
- van Holst, R. J., Lemmens, J. S., Valkenburg, P. M., Peter, J., Veltman, D. J., & Goudriaan, A. E. (2012). Attentional bias and disinhibition toward gaming cues are related to problem gaming in male adolescents. *Journal of Adolescent Health*, *50*, 541-546. doi:10.1016/j.jadohealth.2011.07.006
- Vink, J. M., Beijsterveldt, T. C., Huppertz, C., Bartels, M., & Boomsma, D. I. (2015). Heritability of compulsive Internet use in adolescents. *Addiction Biology*, Epub ahead of print. doi:10.1111/adb.12218
- Volkow, N. D., & Fowler, J. S. (2000). Addiction, a disease of compulsion and drive: Involvement of the orbitofrontal cortex. *Cerebral Cortex*, *10*, 318-325.
- Volkow, N. D., Fowler, J. S., Wang, G. J., & Goldstein, R. Z. (2002). Role of dopamine, the frontal cortex and memory circuits in drug addiction: Insight from imaging studies. *Neurobiology of Learning and Memory*, *78*, 610-624.
- Volkow, N. D., Wang, G. J., Fowler, J. S., & Tomasi, D. (2012). Addiction circuitry in the human brain. *Annual Review of Pharmacology and Toxicology*, *52*, 321-336. doi:10.1146/annurev-pharmtox-010611-134625
- Vollstädt-Klein, S., Löber, S., Kirsch, M., Back, P., Richter, A., Bühler, M., . . . Kiefer, F. (2011). Effects of cue-exposure treatment on neural cue reactivity in alcohol dependence: A randomized trial. *Biological Psychiatry*, *69*, 1060-1066. doi:10.1016/j.biopsych.2010.12.016
- Voon, V., Mole, T. B., Banca, P., Porter, L., Morris, L., Mitchell, S., . . . Irvine, M. (2014). Neural correlates of sexual cue reactivity in individuals with and without compulsive sexual behaviours. *PLoS One*, *9*, e102419. doi:10.1371/journal.pone.0102419
- Wang, C. W., Ho, R. T., Chan, C., L., & Tse, S. (2015). Exploring personality characteristics of Chinese adolescents with internet-related addictive behaviors: Trait differences for gaming addiction and social networking addiction. *Addictive Behaviors*, *42*, 32-35.
- Wang, H., Jin, C., Yuan, K., Shakir, T. M., Mao, C., Niu, X., . . . Zhang, M. (2015). The alteration of gray matter volume and cognitive control in adolescents with internet gaming disorder. *Frontiers in Behavioral Neuroscience*, *9*, 64. doi:10.3389/fnbeh.2015.00064
- Wegmann, E., Stodt, B., & Brand, M. (2015). Addictive use of social networking sites can be explained by the interaction of Internet use expectancies, Internet literacy, and psychopathological symptoms. *Journal of Behavioral Addictions*, *4*, 155-162. doi:10.1556/2006.4.2015.021
- Weinstein, A., Dorani, D., Elhadif, R., Bukovza, Y., Yarmulnik, A., & Dannon, P. (2015). Internet addiction is associated with social anxiety in young adults. *Annals of Clinical Psychiatry*, *27*, 4-9.
- Weinstein, A., & Lejoyeux, M. (2010). Internet addiction or excessive Internet use. *American Journal of Drug and Alcohol Abuse*, *36*, 277-283. doi:10.3109/00952990.2010.491880
- Weinstein, A., & Lejoyeux, M. (2015). New developments on the neurobiological and pharmaco-genetic mechanisms underlying internet and videogame addiction. *American Journal of Addiction*, *24*, 117-125. doi:10.1111/ajad.12110.

- Welberg, L. (2013). Addiction: Craving: A core issue. *Nature Reviews Neuroscience*, *14*, 307. doi:10.1038/nrn3483
- Whang, L. S. M., Lee, S., & Chang, G. (2003). Internet over-users' psychological profiles: A behavior sampling analysis on Internet addiction. *Cyberpsychology & Behavior*, *6*, 143-150. doi:10.1089/109493103321640338
- Widyanto, L., & Griffiths, M. D. (2006). 'Internet Addiction': A critical review. *International Journal of Mental Health and Addiction*, *4*, 31-51. doi:10.1007/s11469-006-9009-9
- Wiers, C. E., Kühn, S., Javadi, A. H., Korucuoglu, O., Wiers, R. W., Walter, H., . . . Bermpohl, F. (2013). Automatic approach bias towards smoking cues is present in smokers but not in ex-smokers. *Psychopharmacology*, *229*, 187-197. doi:10.1007/s00213-013-3098-5
- Wiers, C. E., Ludwig, V. U., Gladwin, T. E., Park, S. Q., Heinz, A., Wiers, R. W., . . . Bermpohl, F. (2015). Effects of cognitive bias modification training on alcohol approach tendencies in male alcohol-dependent patients. *Addiction Biology*, *20*, 990-999. doi:10.1111/adb.12221
- Wiers, R. W., Boelema, S. R., Nikolaou, K., & Gladwin, T. E. (2015). On the development of implicit and control processes in relation to substance use in adolescence. *Current Addiction Reports*, *2*(2), 141-155. doi:10.1007/s40429-015-0053-z
- Wiers, R. W., Eberl, C., Rinck, M., Becker, E. S., & Lindenmeyer, J. (2011). Retraining automatic action tendencies changes alcoholic patients' approach bias for alcohol and improves treatment outcome. *Psychological Science*, *22*(4), 490-497. doi:10.1177/0956797611400615
- Wiers, R. W., & Stacy, A. W. (2006). Implicit cognition and addiction. *Current Directions in Psychological Science*, *15*, 292-296. doi:10.1111/j.1467-8721.2006.00455.x
- Winkler, A., Dörsing, B., Rief, W., Shen, Y., & Glombiewski, J. A. (2013). Treatment of internet addiction: A meta-analysis. *Clinical Psychology Review*, *33*, 317-329. doi:10.1016/j.cpr.2012.12.005
- Wöfling, K., Mörsen, C. P., Duvén, E., Albrecht, U., Grüsser, S. M., & Flor, H. (2011). To gamble or not to gamble: At risk for craving and relapse – learned motivated attention in pathological gambling. *Biological Psychology*, *87*, 275-281.
- Wulfert, E., Maxson, J., & Jardin, B. (2009). Cue-specific reactivity in experienced gamblers. *Psychology of Addictive Behaviors*, *23*, 731-737.
- Xu, Z. C., Turel, O., & Yuan, Y. F. (2012). Online game addiction among adolescents: Motivation and prevention factors. *European Journal of Information Systems*, *21*, 321-340. doi:10.1057/ejis.2011.56
- Yao, Y. W., Wang, L. J., Yip, S. W., Chen, P. R., Li, S., Xu, J., . . . Fang, X. Y. (2015). Impaired decision-making under risk is associated with gaming-specific inhibition deficits among college students with Internet gaming disorder. *Psychiatry Research*, *229*, 302-309. doi:10.1016/j.psychres.2015.07.004
- Yau, Y. H., & Potenza, M. N. (2015). Gambling disorder and other behavioral addictions: Recognition and treatment. *Harvard Review of Psychiatry*, *23*, 134-146. doi:10.1097/HRP.0000000000000051
- Yee, N. (2006). Motivations for play in online games. *Cyberpsychology & Behavior*, *9*, 772-775. doi:10.1089/cpb.2006.9.772
- Yen, J. Y., Yen, C. F., Chen, C. S., Tang, T. C., Huang, T. H., & Ko, C. H. (2011). Cue-induced positive motivational implicit response in young adults with

- Internet gaming addiction. *Psychiatry Research*, 190, 282-286. doi:10.1016/j.psychres.2011.07.003
- Young, K. S. (1996). Addictive use of the Internet: A case that breaks the stereotype. *Psychological Reports*, 79, 899-902. doi:10.2466/pr0.1996.79.3.899
- Young, K. S. (1998). Internet addiction: The emergence of a new clinical disorder. *Cyberpsychology & Behavior*, 3, 237-244. doi:10.1089/cpb.1998.1.237
- Young, K. S. (2004). Internet Addiction: A new clinical phenomenon and its consequences. *American Behavioral Scientist*, 48, 402-415.
- Young, K. S. (2009). Internet addiction: Diagnosis and treatment considerations. *Journal of Contemporary Psychotherapy*, 39, 241-246.
- Young, K. S. (2011). CBT-IA: The first treatment model to address Internet addiction. *Journal of Cognitive Therapy*, 25, 304-312. doi:10.1891/0889-8391.25.4.304
- Young, K. S. (2013). Treatment outcomes using CBT-IA with Internet-addicted patients. *Journal of Behavioral Addictions*, 2, 209-215. doi:10.1556/JBA.2.2013.4.3
- Young, K. S., Pistner, M., O'Mara, J., & Buchanan, J. (1999). Cyber disorders: The mental health concern for the new millennium. *Cyberpsychology & Behavior*, 2, 475-479. doi:10.1089/cpb.1999.2.475
- Young, K. S., Yue, X. D., & Ying, L. (2011). Prevalence estimates and etiologic models of internet addiction. In K. S. Young & C. N. Abreu (Eds.), *Internet addiction* (pp. 3-18). Hoboken, N.J: John Wiley & Sons.
- Yuan, K., Yu, D., Cai, C., Feng, D., Li, Y., Bi, Y., . . . Tian, J. (2016). Frontostriatal circuits, resting state functional connectivity and cognitive control in internet gaming disorder. *Addiction Biology*. doi:10.1111/adb.12348
- Zhang, C., Brook, J. S., Leukefeld, C. G., & Brook, D. W. (2016). Longitudinal psychosocial factors related to symptoms of Internet addiction among adults in early midlife. *Addictive Behaviors*, 62, 65-72. doi:10.1016/j.addbeh.2016.06.019
- Zhou, Y., Lin, F.-C., Du, Y.-S., Qin, L.-D., Zhao, Z.-M., Xu, J.-R., & Lei, H. (2011). Gray matter abnormalities in Internet addiction: A voxel-based morphometry study. *European Journal of Radiology*, 79, 92-95. doi:10.1016/j.ejrad.2009.10.025
- Zhou, Z., Yuan, G., & Yao, J. (2012). Cognitive biases toward Internet game-related pictures and executive deficits in individuals with an Internet game addiction. *PLoS One*, 7, e48961. doi:doi: 10.1371/journal.pone.0048961
- Zhou, Z., Zhu, H., Li, C., & Wang, J. (2014). Internet addictive individuals share impulsivity and executive dysfunction with alcohol-dependent patients. *Frontiers in Behavioral Neuroscience*, 8, 288. doi:10.3389/fnbeh.2014.00288

**Figure 1:** *The model on the development and maintenance of a specific Internet-use disorder. Bold arrows represent the main pathways of the addiction process.*

**Figure 2:** *The reinforcement circle representing a temporal dynamic of the affective and cognitive contributions to the process of development and maintenance of a specific Internet-use disorder. Bold arrows represent the main pathways of the addiction process from the very beginning. The smaller arrows indicate the additional interactions that develop within the addiction process.*

**Figure 3:** *The hypothesized shift from gratification to compensation in the addiction process.*





